

Foliations for solving equations in groups: free, virtually free, and hyperbolic groups.

François Dahmani, Vincent Guirardel

Abstract

We give an algorithm for solving equations and inequations with rational constraints in virtually free groups. Our algorithm is based on Rips' classification of measured band complexes. Using canonical representatives, we deduce an algorithm for solving equations and inequations in all hyperbolic groups (possibly with torsion). Additionally, we can deal with quasi-isometrically embeddable rational constraints.

0 Introduction

0.1 The equations problem

Given a group G , the *equations problem* in G consists in deciding algorithmically whether a system of equations with constants has a solution in G or not. An equation is an equality $w = 1$ where w is a word on a set of variables and their inverses together with constants taken from G . When inequations (i. e. negation of equations) are allowed, we call this problem the *problem of equations and inequations*. In other words, the problem of equations and inequations is equivalent to the decidability of the existential (or universal) theory of G with constants in G .

A solution to the equations problem is quite powerful, as it vastly generalises the word problem, the conjugacy problem, the simultaneous conjugacy problem, etc. When G is abelian, the problem of equations and inequations is easily solved using linear algebra. But already, if G is a free 3-step nilpotent group of rank 2, the equations problem is undecidable ([Tru95], see also [Rom79b]). This is based on Matiyasevich's Theorem saying that one cannot decide the solubility of polynomial equations in $\mathbb{Z}$ [Mat70]. For a free 2-step nilpotent group, the equations problem is solvable if and only if one can decide the solubility of polynomial equations over $\mathbb{Q}$ [Rom79b]. Additionally, if G is a non-commutative free metabelian group, the equations problem in G is unsolvable [Rom79a], but the problem of equations and inequations *without constants* is solvable [Cha95].

The problem of equations and inequations in free groups is natural, and has attracted attention of many people (Lyndon, Appel, Lorents...). The solution of this problem by Makanin in 1982 [Mak82] (with the appropriate correction in [Mak84]) certainly constitutes a milestone in the theory. It has been a source of inspiration for Rips for his study of group actions on $\mathbb{R}$ -trees, and his solution to Morgan and Shalen's conjecture [BF95, GLP94], which found applications in many branches of geometry. It has been a decisive step towards algorithmic, and theoretical description of the set of homomorphisms of a group into a free group (Razborov [Raz84]).

It has also been, together with these developments, a prelude to the far reaching recent solutions to Tarski's problems on the elementary theory of free groups, by Sela [Sel06], and by Kharlampovich and Miasnikov [KM06]. Makanin's algorithm is also the basis of Rips and Sela's solution to the equations problem in torsion free hyperbolic groups that they manage to reduce to the equations problem in a free group [RS95]. Finally, it is crucial in Sela's solution of the isomorphism problem for torsion free hyperbolic groups with finite outer-automorphism group: he makes a delicate use of Makanin's algorithm, and of Rips' classification of actions on $\mathbb{R}$ -trees [Sel95].

Our main result is the following theorem, proved in section 8 (see below for definitions and discussion).

Theorem 1. *There exists an algorithm which takes as input*

- *a presentation of a hyperbolic group G (possibly with torsion),*
- *a finite system of equations and inequations with constants in G , and with quasi-isometrically embeddable rational constraints*

and which decides whether there exists a solution or not.

In a forthcoming paper, we will use this algorithm to give a solution to the isomorphism problem for all hyperbolic groups (possibly with torsion) [DG].

We pursue several goals in this paper. Our first goal is a proof of Theorem 1. As in [RS95, Dah05], our proof is based on canonical representatives, which allow us to reduce to the equations problem in a virtually free group, i. e. a finite extension of a free group.

Our second goal is therefore to give a solution of the equations problem in virtually free groups. This problem easily reduces to a problem of *twisted equations* in a free group.

Last, but not least, we present a new approach to Makanin's algorithm, based on Rips' theory for *foliated band complexes*, allowing us to solve these twisted equations. This occupies the major part of this paper.

We continue this introduction by reviewing different aspects and concepts involved in our strategy.

0.2 Rational constraints

In a group G , the class of *rational subsets* is the smallest class containing finite subsets, and closed under finite union $A \cup B$, product $A \cdot B$, and semi-group generation A^* . Equivalently, if S is a finite generating set of G , a subset $A \subset G$ is rational if it is the image in G of a regular language L of the free monoid on $S \cup S^{-1}$, a regular language being a language recognised by a finite state automaton. Solving a system of equations with *rational constraints* consists in solving this system of equations with the requirement that each variable x lies in a rational subset $\mathcal{R}_x$ given in advance.

In a hyperbolic group, we propose the class of *quasi-isometrically embeddable* rational subsets as the suitable class for constraints. A rational subset $A \subset G$ is quasi-isometrically embeddable if one can choose the regular language L in the free monoid to consist of quasigeodesics (with uniform constants). For example, a quasi-convex subgroup H of a hyperbolic group G is a quasi-isometrically embeddable

rational subset. Note that by [Kap96], one can compute a finite state automaton representing H from a finite generating set. In general, the class of rational subsets of a group is not closed under complementation or intersection. However, the set of quasi-isometrically embeddable rational subsets of a hyperbolic group is a Boolean algebra (Cor. 9.6). For instance, the complement of a quasi-convex subgroup is also a quasi-isometrically embeddable rational subset. In particular, inequations in a hyperbolic group can be encoded using quasi-isometrically embeddable rational constraints of the form $\mathcal{R} = G \setminus \{1\}$. In a virtually free group, every rational subset is quasi-isometrically embeddable, and the set of all rational subsets is a Boolean algebra.

Equations with rational constraints in a free monoid were first considered by Schulz [Sch92]. Following an approach of Plandowski for free monoids, Diekert, Gutierrez and Hagenah proved that systems of equations, inequations, and rational constraints in free groups are algorithmically solvable [Pla04, DGH05].

The use of rational constraints in systems of equations turns out to be rather powerful. The problem of equations and inequations for right angled Artin groups has been reduced by Diekert and Muscholl to a problem of equations with rational constraints in free groups [DM06]. This has been generalised by Diekert and Lohrey to free products, direct products, and graph products of certain groups [DL08]. It is the key tool to extend Rips and Sela's solution to the equations problem for torsion free hyperbolic groups into a solution to the problem of equations and inequations [Dah05]. It greatly streamlines the solution of the isomorphism problem for torsion free hyperbolic groups, and allows substantial generalisations [DG08, DG]. It plays an important role in our solution to the equations problem in virtually free groups (Th.3 below), and in Lohrey and Senizergues' independent solution [LS06a], even if the initial problem does not involve inequations or rational constraints.

For the sake of illustration, let us present two elementary applications of the use of rational constraints in a system of equations. Let F_2 be a free group of rank 2. Given a finitely generated subgroup $H < F_2$ (or more generally any rational subset), and an element $x \in F_2$, one can decide if there is an automorphism of F_2 sending x into H . Indeed, consider a, b a basis of F_2 , and write $x = w(a, b)$ as a word on a, b . The orbit of x under $\text{Aut}(F_2)$ intersects H if and only if there exists a basis u, v of F_2 such that $w(u, v) \in H$. By a theorem of Dehn, Magnus and Nielsen, (u, v) is a basis if and only if $\exists g, [u, v]^g = [a, b]^{\pm 1}$. Therefore, the orbit of x intersects H if and only if the system of equations with rational constraints

$$\begin{cases} [u, v]^g = [a, b]^{\pm 1} \\ z = w(u, v), \quad z \in H \end{cases}$$

in the variables u, v, g, z has a solution.

Our second application is an immediate consequence of Theorem 1:

Corollary 2. *Let G be a hyperbolic group. Given a quasiconvex subgroup $H < G$, one can decide its malnormality by solving the system $z = yxy^{-1}$, $z \neq 1$ with rational constraints $z, x \in H$, $y \notin H$.*

In presence of torsion, almost malnormality (meaning that $yHy^{-1} \cap H$ is finite when $y \notin H$) can be checked similarly by replacing the inequality $z \neq 1$ by an inequality $z^{N!} \neq 1$ where N is a bound on the order of torsion in G .

This result does not seem to appear in the literature. This is a variant of problem H14 in [BMS] and of Question 3 in [BW01]. Without assumption of quasiconvexity, malnormality is undecidable [BW01].

0.3 Lifting equations and rational constraints to a virtually free group

Following the strategy initiated in [RS95], and continued in [Dah05], we now explain how to reduce Theorem 1 to the problem of solving equations with rational constraints in a virtually free group (see Section 9).

In [RS95], Rips and Sela introduced *canonical representatives* for torsion free hyperbolic groups, which enabled them to reduce the equations problem in a torsion free hyperbolic group to the equations problem in a free group. These canonical representatives are paths in the Cayley graph of G which satisfy some *path equations* representing the initial equations. Such paths correspond to words on the generating system of G , and thus to elements of the corresponding free group. In presence of torsion, canonical representatives need to be interpreted as paths in a barycentric subdivision X of a Rips complex of G . The action of G on X is not free in general. The quotient of the 1-skeleton $X^{(1)}/G$ is a finite graph of finite groups, whose fundamental group is a virtually free group V . Path equations in X are then interpreted in terms of equations in V .

Another task that needs to be done, is to lift rational constraints to V . This can be done because both canonical representatives and paths representing the rational subsets are quasi-geodesics. This part of the argument is similar to Cannon's argument showing that the language of geodesics is a regular language [Can84, CDP90].

This allows to reduce Theorem 1 to the following result (proved in section 8):

Theorem 3. *The problem of equations with rational constraints is solvable in finite extensions of free groups.*

More precisely, there exists an algorithm which takes as input a presentation of a virtually free group G , and a system of equations and inequations with constants in G , together with a set of rational constraints, and which decides whether there exists a solution or not.

This result was independently obtained in a more general form in [LS06a]. Unlike [LS06a], our approach does not rely on an existing solution of the equations problem in free groups. Instead, we give a new proof, based on Rips' theory for foliated band complexes.

0.4 Twisted equations and virtually free groups

A *twisted equation* in a group G is an equation of the form $\varphi_1(x_1) \dots \varphi_n(x_n) = 1$ where each φ_i is a fixed automorphism in $\text{Aut}(G)$, and x_i is a variable or a constant. For instance, twisted conjugacy involves a simple example of a twisted equation: given an automorphism $\varphi \in \text{Aut}(G)$, two elements $a, b \in G$ are twisted conjugate if there exists $x \in G$ such that $xa\varphi(x^{-1}) = b$.

When one considers equations in a finite extension $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$ (Q is finite), twisted equations appear naturally. Indeed one can replace an equation $xyz = 1$, by a disjunction of equations $(\tilde{q}_x n_x)(\tilde{q}_y n_y)(\tilde{q}_z n_z) = 1$, where the constants

$\tilde{q}_x, \tilde{q}_y, \tilde{q}_z$ are chosen in a given cross-section of Q , and n_x, n_y, n_z are new unknowns in N . Gathering the elements $\tilde{q}_x, \tilde{q}_y, \tilde{q}_z$ to the left amounts to twist n_x and n_y by suitable automorphisms. Thus, the solubility of equations in G reduces to the solubility of finitely many systems of twisted equations in N . The twisting morphisms occurring in this manner are quite particular because they generate a finite subgroup of the outer automorphism group $\text{Out}(N)$.

In the Kurovka notebook, Makanin asked about the problem of twisted equations [MK02, Problem 10.26(b)]. We are able to give a positive answer to Makanin's question above assuming that the given automorphisms generate a finite subgroup of $\text{Out}(F)$:

Theorem 4. *There exists an algorithm that takes as input a basis of a free group F , a finite set Φ of automorphisms of F whose image in $\text{Out}(F)$ generate a finite subgroup, and a system of twisted equations with rational constraints in F (with twisting automorphisms in Φ) and that decides whether there is a solution or not.*

As explained above, Theorem 3 easily follows from Theorem 4 (see Section 2). Moreover, we give a trick to reduce to the case where the twisting automorphisms of F permute the elements of $S \cup S^{-1}$ for some free basis S of F . This trick is related to the Zimmerman-Culler Theorem [Zim81, Cul84] which realises any finite subgroup of $\text{Out}(F)$ as a finite group H of automorphisms of a graph X whose fundamental group is F . Because H may fail to fix a point in X , H may fail to lift to a finite subgroup of $\text{Aut}(S)$. This is why we need to embed F into a larger free group $\hat{F}$, whose basis is the set of edges of X . The group H is then realised as a subgroup of $\text{Aut}(\hat{F})$ permuting the basis elements, and H preserves the conjugacy class of F . The initial system of equations gives a new system of equations in $\hat{F}$, and we add rational constraints saying that the variables should live in F . This reduction is the content of Proposition 2.4 in the broader context of equations with rational constraints.

0.5 Dynamical and geometric aspects

We now focus on the equations problem in a free group. Makanin and Razborov developed a combinatorial machinery to encode equality of subwords occurring in a solution of an equation. An interesting and well written account on Makanin's algorithm for equations in free monoids (a simplified version of the case of free groups, that does not imply a solution for free groups) was given in [Die02], and another one on Makanin and Razborov's algorithm for free groups was given in [KM05]. As we said, Rips was inspired by this machinery for his study of foliated band complexes, whose dynamics, on the other hand, reflect actions on $\mathbb{R}$ -trees. Our strategy is to reverse this flow of ideas, and use Rips' classification of foliated band complexes to prove that the algorithm we propose always stops.

We hope that this point of view on Makanin's algorithm will be of interest to an audience concerned with equations problems and also to an audience concerned with geometry and dynamics of group actions.

Rips' theory is an understanding of actions of finitely generated groups on $\mathbb{R}$ -trees. Recall that an $\mathbb{R}$ -tree is a geodesic metric space in which any two points are joined by a unique injective path. Following Sela, let us try to explain how the equations problem in a free group F is related to $\mathbb{R}$ -trees [Sel01].

The equations problem in F is about homomorphisms of finitely presented groups to F . Indeed, a system of equations on a set of unknowns X is a finite set $\mathcal{E}$ of words in the free product $F * \langle X \rangle$, and a solution of this system of equations corresponds to a morphism from $G_{\mathcal{E}} = F * \langle X \rangle / \langle\langle \mathcal{E} \rangle\rangle$ to F which is the identity on F . Each such morphism gives rise to an action of $G_{\mathcal{E}}$ on the Cayley graph of F , a simplicial tree. One can rescale this tree in order to normalise the maximal displacement of the generators of $G_{\mathcal{E}}$. For an infinite sequence of solutions, the corresponding actions of $G_{\mathcal{E}}$ on the Cayley graph converge to an action of $G_{\mathcal{E}}$ on some $\mathbb{R}$ -tree.

Rips' theory says that under suitable hypotheses, this action can be understood in terms of actions on simplicial trees, and actions on $\mathbb{R}$ -trees dual to minimal measured foliations on 2-complexes. The main result of Rips' theory is a classification of those minimal measured foliations into three types:

- homogeneous type (also known as axial or toral), whose dual $\mathbb{R}$ -tree is a line
- surface type (also known as interval exchange): the $\mathbb{R}$ -tree is dual to a measured foliation on a surface (or a 2-orbifold)
- exotic type (also known as Levitt or thin).

One can try to use these ideas to decide whether a given system of equations has a solution, and to look for a *shortest* solution (e.g. in terms of the maximal displacement of the generators). If we have some solutions such that the corresponding actions of $G_{\mathcal{E}}$ on the Cayley graph of F are close enough to the limiting $\mathbb{R}$ -tree, we can apply Sela's *shortening argument*. This argument says that, under suitable hypotheses, there is a quotient of $G_{\mathcal{E}}$ through which the actions factorise, and automorphisms of this quotient that *shorten* all nearby solutions. These solutions are not shortest and can therefore be ignored. Using some compactness argument, only finitely many solutions do not lie in such a shortening neighbourhood of a limiting $\mathbb{R}$ -tree. One can hope to bound the length of these remaining solutions and check by hand if such a solution exists.




Figure 1: A prelamination is a finite disjoint union of *leaf segments*.

A major problem in this approach is that we cannot even start as we don't know if our system of equations has any solution (this is what we have to decide). Therefore, we cannot work with *actual* solutions. Instead, we work with *potential* solutions. An actual solution, i. e. a morphism $G_{\mathcal{E}} \rightarrow F$, can be represented by a continuous map from a presentation complex X of $G_{\mathcal{E}}$ to a wedge of circles Y . The preimage of midpoints of edges of Y is a combinatorial *lamination* of X . Instead of working with laminations, we work with *prelaminations* that play the role of potential solutions. A subcomplex of a lamination is a typical example of a prelamination,

but prelaminations are defined by local conditions and are not required to extend to an actual lamination (See Figure 1).

Our proof shows that when a prelamination is very long, then any actual solution corresponding to this prelamination can be shortened, and therefore ignored. This is the key of the termination of our algorithm.

0.6 Overview of the main algorithm

Let us describe the approach to solving equations in free groups developed in this paper.

The first step toward Theorem 4 is to encode a system of triangulated equations in a *band complex*. Band complexes are Makanin’s *generalised equations*, but they also play the role of a presentation complex of $G_{\mathcal{E}}$ in the above discussion. A band complex consists of D a disjoint union of non-degenerate segments (one segment for each variable or constant involved in $\mathcal{E}$) together with a finite set of rectangles (called *bands*) attached to D by two opposite sides called its bases. If one considers twisted equations in the context described above, then each band carries a specific automorphism. If one considers rational constraints, then subsegments of D carry regular languages.

A *solution* of the band complex in the free group $\langle S \rangle$ is a labelling of D by reduced words on $S \cup S^{-1}$ so that both bases of each band are labelled by the same reduced word (up to composing by the automorphism of the band), and so that the label of each subsegment satisfies the associated rational constraint. For instance, to encode constants, one can use regular languages consisting of a single word to impose the labelling on some subsets of D . Solving twisted equations in free groups easily reduces to deciding the existence of a solution to such a band complex (see Section 3). To simplify this presentation, we now forget about rational constraints.

Our *main algorithm* 8.2 decides whether a given band complex admits a solution (i. e. a labelling as above) using prelaminations. A prelamination is a finite disjoint union of *leaf segments*, and a leaf segment is a segment contained in a band and joining its two bases (see Figure 1). A prelamination is *induced* by a solution if its leaf segments join matching subwords of the labelling. A prelamination is *complete* if one cannot extend any leaf (i. e. if it is an actual lamination). Given a complete prelamination, it is easy to decide if there exists a solution that induces it.

We can explore the space of possible prelaminations, in quest of a complete prelamination induced by a shortest solution. The main concern is how to detect that there is no solution.

For each complete prelamination, we can decide if it is induced by a solution or not. If it is, we are done, and otherwise, we reject this prelamination. For each incomplete prelamination $\mathcal{L}$, we run a *prelamination analyser* which tries to find a certificate ensuring that no shortest solution can induce $\mathcal{L}$. In this case one can reject $\mathcal{L}$. The analyser may fail to find any such certificate and might say “I don’t know”. There are several kinds of certificates: detection of an incompatibility with constants, non-existence of an invariant (combinatorial) transverse measure, or a *shortening certificate* proving that if some solution induces $\mathcal{L}$, then there is a shorter solution.

Note that if a lamination $\mathcal{L}'$ extends a rejected $\mathcal{L}$, then $\mathcal{L}'$ cannot be induced by a shortest solution. If all prelaminations remaining to analyse are extensions of rejected ones, we know that there is no shortest solution, hence no solution at all,

and the machine stops.

To prove that this algorithm works, we assume that there is no solution, and show that the algorithm rejects all sufficiently long prelaminations by producing appropriate shortening certificates.

This is where we use Rips' classification of measured foliations on band complexes. Assume by contradiction that the algorithm does not stop. By extracting a limit of an increasing sequence of non rejected prelaminations $\mathcal{L}_1 \subset \mathcal{L}_2 \subset \dots$, one can construct a topological foliation on the band complex with an invariant measure. We prove that this measure has no atom. Assume for simplicity that it has full support (this is not true in general). One can decompose the foliated band complex into minimal components. By Rips' theory, they are classified as homogeneous, exotic, and surface type components.

In presence of a homogeneous component, for all i large enough, $\mathcal{L}_i$ shows large repetition patterns that force solutions inducing $\mathcal{L}_i$ to have subwords that are arbitrarily high powers. By Bulitko's Lemma, there is a computable a priori bound on such powers in a shortest solution. One can therefore produce a shortening certificate for i large enough.

In presence of a surface or exotic component, one can perform, as in the Rips Machine, an infinite sequence of moves on the foliated band complex Σ . These moves don't increase the complexity of Σ , so finitely many unfoliated band complexes are visited. Assume for simplicity that Σ itself is visited twice. For i large enough, the corresponding moves are compatible with $\mathcal{L}_i$, and transform any solution of Σ inducing $\mathcal{L}_i$ into a shorter solution of Σ . These moves provide a shortening certificate for $\mathcal{L}_i$: if $\mathcal{L}_i$ is induced by some solution, then this solution cannot be shortest. When Σ itself is not visited twice, this argument needs to be refined, and shortening must be performed by at least a certain Lipschitz factor.

The running time of our algorithm does not seem to be very good. Makanin's algorithm (in its corrected version [Mak84]) is known not to be primitive recursive [KP98], and we don't see why ours should be better. Using some data compression on words, Plandowski proposed an algorithm of much better complexity (polynomial in space) [Pla04, DGH05].

0.7 Organisation of the paper

Sections 1 to 3 set up the vocabulary, and reduce the equations problem in virtually free groups to band complexes. Section 4 is about Bulitko's Lemma on the periodicity exponent of minimal solutions. Section 5 introduces prelaminations and related notions. Section 6 describes the prelamination generator and the prelamination analyser, and gives a precise definition of a *shortening sequence of moves* that we use as a certificate for rejection. Section 7 contains the main part of the argument. Limits of sequences of prelaminations are analysed using Rips' theory, and existence of shortening sequences of moves is proved. Section 8 contains a detailed account on the main algorithm. This section can be read independently, admitting a certain number of well identified results of the previous sections. Although based on Theorem 3, Section 9 is independent from the rest of the paper. It deals with the equations problem in hyperbolic groups using canonical representatives, and with quasi-isometrically embeddable rational constraints.

In a first reading, one can first forget about rational constraints. Moreover, if one is only interested in the case of free groups, i. e. with untwisted equations, one

can safely ignore all technical considerations about Möbius strips (see Remark 5.3).

Contents

0	Introduction	1
0.1	The equations problem	1
0.2	Rational constraints	2
0.3	Lifting equations and rational constraints to a virtually free group	4
0.4	Twisted equations and virtually free groups	4
0.5	Dynamical and geometric aspects	5
0.6	Overview of the main algorithm	7
0.7	Organisation of the paper	8
1	Preliminaries	10
1.1	Regular languages	10
1.2	Rational subsets	11
1.3	Rational subsets of a virtually free group	12
2	Equations and twisted equations	13
2.1	Reduction to triangular equations	14
2.2	Twisted equations	14
3	Band complexes	17
3.1	Band complexes	17
3.2	Solution of a band complex	17
3.3	Rational constraints	18
3.4	From twisted equations to band complexes	18
3.5	Standard forms of rational constraints	20
4	Bulitko's Lemma: bounding the exponent of periodicity	22
4.1	Normal decomposition with respect to powers	22
4.2	Special segments and concatenation	23
4.3	Proof of Bulitko's Lemma	23
5	Prelaminations	25
5.1	Definition	25
5.2	Möbius strips	26
5.3	Rational constraints on the prelamination	27
5.4	Prelaminations and rational constraints induced by a solution	27
6	Algorithms on prelaminations	28
6.1	A first algorithm: the prelamination generator	28
6.2	A second algorithm: the prelamination analyser	29
7	Asymptotic of prelaminations: Rips band complexes	35
7.1	Limit measured foliation	36
7.2	Moves	41
7.3	Minimal components	44
7.4	Homogeneous components	48
7.5	Independent bands	48
7.6	The pruning process	51

8	Endgame	54
8.1	Finding solutions of band complexes	54
8.2	Reduction of main theorems to band complexes	57
9	Equations in hyperbolic groups with torsion	57
9.1	Quasi-isometrically embeddable rational subsets	58
9.2	Full sublanguage of a set of quasigeodesics	59
9.3	Canonical representatives in hyperbolic groups with torsion	62
9.4	Lifting equations to V	65
9.5	Lifting rational constraints to V , and proof of the theorem	66

1 Preliminaries

1.1 Regular languages

Let S be a finite set, $\overline{S}$ a copy of S , and $S_{\pm} = S \sqcup \overline{S}$ endowed with the canonical involution $s \mapsto \overline{s}$ exchanging S and $\overline{S}$. Let $S_{\pm}^*$ be the free monoid on $S_{\pm}$, endowed with the involution $w = s_1 \dots s_n \mapsto \overline{w} = \overline{s}_n \dots \overline{s}_1$.

An *automaton* over $S_{\pm}$ is a directed graph where each (oriented) edge is labelled by some element of $S_{\pm}$ together with two finite subsets $Start, Accept$ of the set of vertices of this graph. The *language* $\mathcal{R}_A \subset S_{\pm}^*$ accepted by an automaton A is the set of words labelled by directed paths starting from a vertex in $Start$ and finishing at a vertex in $Accept$. A subset $\mathcal{R} \subset S_{\pm}^*$ is a *regular language* if $\mathcal{R} = \mathcal{R}_A$ for some automaton A .

If $\mathcal{R}_1, \mathcal{R}_2$ are regular languages, then so are $\mathcal{R}_1 \cup \mathcal{R}_2$, $\mathcal{R}_1 \cdot \mathcal{R}_2$ (the product set), and $\mathcal{R}_1^*$ (the submonoid generated by $\mathcal{R}_1$). Kleene's theorem asserts that the class of regular languages is the smallest class containing finite subsets, and closed under these operations.

The class of regular languages is actually a Boolean algebra. Moreover, one can algorithmically compute automata for $\mathcal{R}_1 \cup \mathcal{R}_2$, $\mathcal{R}_1 \cap \mathcal{R}_2$, $S_{\pm}^* \setminus \mathcal{R}_1$, $\mathcal{R}_1 \cdot \mathcal{R}_2$ and $\mathcal{R}_1^*$ from automata representing $\mathcal{R}_1$ and $\mathcal{R}_2$. Similarly, if $\mathcal{R}$ is rational, then so is its image $\overline{\mathcal{R}}$ under the involution $w \mapsto \overline{w}$, and the corresponding automaton can be computed.

Example 1.1. Assume that $\pi : S_{\pm}^* \rightarrow F$ is a morphism to a finite group F , and consider a subset $E \subset F$. Let A be the Cayley graph of F relative to $\pi(S)$.

Then $\pi^{-1}(E)$ is a regular language, corresponding to the automaton A where $Start = \{1\}$ and $Accept = E$.

Not all regular languages are of this form but this is true if one replaces the finite group F by a finite monoid. Let's be more specific. Denote by $\mathcal{M}_n$ the finite monoid of matrices with Boolean entries in $\{0, 1\} = \{\mathbf{true}, \mathbf{false}\}$, where the product of $C = AB$ of $A = (a_{ij})$ and $B = (b_{jk})$ is $(c_{ik}) = (\bigvee_{j=1}^n a_{ij} \wedge b_{jk})$ and $\vee, \wedge$ denote **or** and **and**. Consider an automaton A over $S_{\pm}$ with vertices $v_1, \dots, v_n$. To each $s \in S_{\pm}$ corresponds the matrix M_s whose entry (i, j) is 1 if and only if there is an edge labelled s from v_i to v_j . The assignment $s \mapsto M_s$ extends to a morphism $\rho : S_{\pm}^* \rightarrow \mathcal{M}_n$ and the language $\mathcal{R}_A$ accepted by A is the preimage under ρ of the set of matrices μ having a non-zero entry (i, j) where $v_i \in Start$ and $v_j \in Accept$. Conversely, given a morphism $\rho : S_{\pm}^* \rightarrow \mathcal{M}$ to a finite monoid $\mathcal{M}$ and a subset $\mu \subset \mathcal{M}$, $\rho^{-1}(\mu)$ is a regular language corresponding to the following automaton:

its vertex set is $\mathcal{M}$, its edges labelled by s are given by the multiplication by $\rho(s)$, $Start = \{\text{id}\}$ and $Accept = \mu$. Clearly, one can compute ρ and μ from an automaton, and conversely.

Remark 1.2. This (well known) construction implies the (well known) fact that one can make an automaton deterministic, and in particular, assume that $Start$ is a single vertex of the automaton.

1.2 Rational subsets

Consider a group G , S a finite generating set, and $\pi : S_{\pm}^* \rightarrow G$ the natural morphism mapping $S \cup \overline{S}$ to $S \cup S^{-1}$.

Definition 1.3. A rational subset of a group G is the image under π of a regular language of $S_{\pm}^*$.

Equivalently, the class of rational subsets of G is the smallest class containing finite subsets, and closed under union, product, and submonoid generation.

The equivalence between the two definitions follows from Kleene's theorem. In particular, the notion of rational subset does not depend on the choice of the generating set S . We say that a rational subset $\mathcal{R} \subset G$ is *represented* by an automaton A over $S_{\pm}$ when $\pi(\mathcal{R}_A) = \mathcal{R}$.

Note that if $\mathcal{R}$ is rational, so is $\mathcal{R}^{-1}$. Although the class of regular languages of $S_{\pm}^*$ is a Boolean algebra, the class of rational subsets of a group G is not closed under intersection or complementation in general. Indeed, it is easy to see that subgroups are rational subsets if and only if they are finitely generated, but it can happen that the intersection of two such subgroups is not finitely generated.

Lemma 1.4. Let $f : G \rightarrow G'$ be a morphism, S, S' be finite generating sets of G, G' , and A an automaton on S representing some rational subset $\mathcal{R} \subset G$.

Then $f(\mathcal{R})$ is a rational subset of G' , and knowing an expression of $f(s)$ as an S' -word, one can compute an automaton for $f(\mathcal{R})$.

Proof. For each $s \in S \cup \overline{S}$, write $f(s) = s'_1 \dots s'_n$ where $s'_i \in S' \cup \overline{S'}$. Replace each directed edge of A labelled s by a directed segment of length n labelled $s'_1 \dots s'_n$. The automaton obtained in this way clearly represents $f(\mathcal{R})$. $\square$

The following lemma follows immediately from the analogous fact concerning regular languages of $S_{\pm}^*$. It holds without assumption on the group G .

Lemma 1.5. Given two automata A_1, A_2 defining some rational subsets $\mathcal{R}_1, \mathcal{R}_2$ of G , one can algorithmically compute some automata representing $\mathcal{R}_1 \cup \mathcal{R}_2$, $\mathcal{R}_1 \cdot \mathcal{R}_2$, $\mathcal{R}_1^*$ and $\mathcal{R}_1^{-1}$.

When G is a free group, or even a virtually free group, the class of rational subsets is a Boolean algebra (see below). For the free group, this is based on the following fact. We denote by $\langle S \rangle$ the free group with free basis S .

Lemma 1.6 ([Ber79, Proposition 2.8 p59]). *For any rational subset $\mathcal{R}$ in the free group $\langle S \rangle$, consider the set $\tilde{\mathcal{R}} \subset S_{\pm}^*$ of reduced words representing elements of $\mathcal{R}$.*

Then $\tilde{\mathcal{R}}$ is a regular language, and an automaton representing $\tilde{\mathcal{R}}$ can be computed from an automaton representing $\mathcal{R}$.

It follows that the class of rational subsets of a free group is a Boolean algebra: denoting by Red the regular language of reduced words of $S_\pm^*$, one has $\pi(\text{Red} \setminus \tilde{\mathcal{R}}) = \langle S \rangle \setminus \mathcal{R}$ and $\pi(\tilde{\mathcal{R}}_1 \cap \tilde{\mathcal{R}}_2) = \mathcal{R}_1 \cap \mathcal{R}_2$. Moreover, if S is a basis of the free group F , automata over $S_\pm$ representing the obtained rational subsets can be computed from automata over $S_\pm$ for the initial ones.

1.3 Rational subsets of a virtually free group

When G is virtually free, the fact that rational subsets form a Boolean algebra follows from recent work by Lohrey and Senizergues [LS06b] (their result is more general but quite complicated). We propose here a short proof of what we need.

Remark 1.7. From a presentation of a virtually free group V , one can compute a free basis of a normal finite index free subgroup F and a representative $g_a \in V$ of each element $a \in V/F$. Indeed, by the Reidemeister-Schreier process, one can enumerate all normal finite index subgroups of V , with presentations, and by Tietze transformations, all presentations of these subgroups. One will find one that is a presentation with no relator, i. e. a free basis of a free group F . Then the finite group V/F and lifts in V are easy to compute.

Lemma 1.8. *Let V be a virtually free group given by some presentation with generating set S_V . Let $F \triangleleft V$ be a normal free subgroup of finite index, given by a free basis S , expressed as a set of words on S_V . Let $\mathcal{R}$ be a rational subset of F (hence of V).*

Given an automaton over $(S_V \cup \overline{S}_V)$ representing $\mathcal{R}$, one can compute an automaton over $S_\pm$ representing it, and vice-versa.

Proof. Computing an automaton over $(S_V \cup \overline{S}_V)$ from one over $S_\pm$ simply follows from Lemma 1.4 applied to the embedding $F \rightarrow V$.

Let $\pi : (S_V \cup \overline{S}_V)^* \rightarrow V$ be the natural map. Let A be an automaton over $S_V \cup \overline{S}_V$ accepting a language $\tilde{\mathcal{R}} \subset (S_V \cup \overline{S}_V)^*$ with $\pi(\tilde{\mathcal{R}}) = \mathcal{R}$. Let V_A be its set of vertices, and assume that Start is a single vertex (this can be assumed by Remark 1.2). Denote by $\rho : (S_V \cup \overline{S}_V)^* \rightarrow V/F$ the composition of π with the quotient map.

We construct a new automaton A' with vertex set $V_{A'} = V_A \times V/F$. We put an edge labelled $u \in S_V \cup \overline{S}_V$ from (v_1, a) to (v_2, b) if there is an edge labelled u from v_1 to v_2 in A , and $b = a\rho(u)$. We take $\text{Start}' = (\text{Start}, 1)$ and $\text{Accept}' = \text{Accept} \times \{1\}$. Since $\mathcal{R} \subset F$, the language accepted by A' is precisely $\tilde{\mathcal{R}}$.

For each $a \in V/F$ choose a representative $g_a \in V$ with $g_a = 1_V$ for $a = 1$. We also think of g_a as a word on $(S_V \cup \overline{S}_V)$, and we note that such g_a can be computed. We construct an automaton A'' over $S_\pm$ as follows. Consider an edge e labelled u and joining (v_1, a) to (v_2, b) . Note that $u' = g_a u g_b^{-1} \in F$, and write $u' = s_1 \dots s_k$ as a word on $S_\pm$ (this can be done algorithmically). Replace the edge e by a directed segment labelled by $s_1 \dots s_k$. Do this for every edge of A' , take $\text{Start}'' = \text{Start}'$ and $\text{Accept}'' = \text{Accept}'$, and call A'' the obtained automaton.

We claim that the language $\tilde{\mathcal{R}}''$ accepted by A'' satisfies $\pi(\tilde{\mathcal{R}}'') = \mathcal{R}$. Indeed, assume that $u_1 \dots u_n \in (S_V \cup \overline{S}_V)^*$ is accepted by A' , and consider $a_i = \rho(u_1 \dots u_i)$. One has $a_0 = 1$ since the path starts at Start' , and $a_n = 1$ since $\pi(u_1 \dots u_n) \in F$. Then the $S_\pm$ -word $u'_1 \dots u'_n = (1u_1g_{a_1}^{-1})(g_{a_1}u_2g_{a_2}^{-1}) \dots (g_{a_{n-1}}u_n1)$ is accepted by A'' , and has the same image under π as $u_1 \dots u_n$. Similarly any word accepted by A'' maps to $\mathcal{R}$. $\square$

Lemma 1.9. *Let V be a virtually free group, and $F \triangleleft V$ be a normal free subgroup of finite index, and let $g_1, \dots, g_k$ be representatives of the left cosets of F in V .*

A subset $\mathcal{R} \subset V$ is rational if and only if for all i , $(g_i^{-1}\mathcal{R}) \cap F$ is rational in F .

Let S_V be a generating set for V , and S a basis of F . Given automata over $S_{\pm}$ representing each $(g_i^{-1}\mathcal{R}) \cap F$, one can explicitly compute an automaton over $S_V \cup \overline{S}_V$ representing $\mathcal{R}$, and conversely, given an automaton over $S_V \cup \overline{S}_V$ representing $\mathcal{R}$, and $g_1, \dots, g_k$, one can compute an automaton over $S_{\pm}$ representing $(g_i^{-1}\mathcal{R}) \cap F$.

Proof. If $(g_i^{-1}\mathcal{R}) \cap F$ is a rational subset of F , it is rational in V . Therefore, the sets $\mathcal{R} \cap g_i F$ are rational in V , and so is their union. Given an automaton over $S_{\pm}$ representing $(g_i^{-1}\mathcal{R}) \cap F$, one can compute an automaton over $S_V \cup \overline{S}_V$ representing it by Lemma 1.8, hence an automaton over $S_V \cup \overline{S}_V$ representing $\mathcal{R}$ by Lemma 1.5.

Conversely, assume that $\mathcal{R}$ is rational in V . Denote by $\pi : (S_V \cup \overline{S}_V)^* \twoheadrightarrow V$ the natural morphism, and $\tilde{\mathcal{R}} \subset (S_V \cup \overline{S}_V)^*$ a rational language such that $\mathcal{R} = \pi(\tilde{\mathcal{R}})$. By Example 1.1, $\pi^{-1}(F)$ is a regular language, and so is $\tilde{\mathcal{R}} \cap \pi^{-1}(F)$. Thus $\mathcal{R} \cap F$ is rational, and one can compute an automaton A over $S_V \cup \overline{S}_V$ representing it. By Lemma 1.8, we can turn it into an automaton over $S_{\pm}$. $\square$

Lemma 1.10. *Let V be a virtually free group, and $\mathcal{R}, \mathcal{R}'$ be rational subsets.*

Then $V \setminus \mathcal{R}$ and $\mathcal{R} \cap \mathcal{R}'$ are rational, and one can compute automata representing them from automata representing $\mathcal{R}$ and $\mathcal{R}'$.

Proof. By Lemma 1.9, one can write $\mathcal{R} = g_1\mathcal{L}_1 \sqcup \dots \sqcup g_k\mathcal{L}_k$ with $\mathcal{L}_i$ rational in F . Then $V \setminus \mathcal{R} = g_1(F \setminus \mathcal{L}_1) \sqcup \dots \sqcup g_k(F \setminus \mathcal{L}_k)$ is rational since in the free group F , $(F \setminus \mathcal{L}_i)$ is rational. Moreover, one can compute automata for $\mathcal{L}_i$ by Lemma 1.9, hence for $F \setminus \mathcal{L}_i$, and for $V \setminus \mathcal{R}$.

For the second assertion, simply write $\mathcal{R} \cap \mathcal{R}' = (\mathcal{R}^c \cup \mathcal{R}'^c)^c$. $\square$

2 Equations and twisted equations

In this section, we start by giving a formal definition of a system of equations with rational constraints in a group G . Then we explain how to translate a system of equations in a virtually free group into a disjunction of systems of twisted equations in a free group.

Definition 2.1 (Equations with rational constraints). *Consider a finite set of variables X and $X_{\pm} = X \sqcup \overline{X}$ with the natural involution $x \mapsto \overline{x}$.*

A system of equations $\mathcal{E}$ is a finite set of words $w = x_1 \dots x_n$ in $X_{\pm}^$ (representing the equation $x_1 \dots x_n = 1$).*

A set of rational constraints for $\mathcal{E}$ is a tuple $\underline{\mathcal{R}} = (\mathcal{R}_x)_{x \in X}$ where for each $x \in X$, $\mathcal{R}_x \subset G$ is a rational subset.

A solution of $(\mathcal{E}, \underline{\mathcal{R}})$ is a tuple $\underline{g} = (g_x)_{x \in X} \in G^X$ with $g_x \in \mathcal{R}_x$ for each $x \in X$, and such that for each $w = x_1 \dots x_n \in \mathcal{E}$, $g_{x_1} \dots g_{x_n} = 1$ in G where we define $g_{\overline{x}} = g_x^{-1}$ for each $x \in X$.

Abusing notations, if $(g_x)_{x \in X}$ is a solution of $\mathcal{E}$, we will also call solution the corresponding tuple $(g_x)_{x \in X \cup \overline{X}}$ where $g_{\overline{x}} = g_x^{-1}$ for each $x \in X$.

Constants in a system of equations can be encoded with rational constraints: a constant $g_0 \in G$ is a variable x with corresponding constraint $\mathcal{R}_x = \{g_0\}$. If $G \setminus \{1\}$ is

a rational subset, then inequations can be encoded with rational constraints: replace each inequation $w \neq 1$ by an equation $w = y$ where y is a new variable with rational constraint $\mathcal{R}_y = G \setminus \{1\}$.

2.1 Reduction to triangular equations

We fix V a finitely generated virtually free group. Classically, a system of equations with rational constraints is equivalent to a triangular system (i. e. in which every $w \in \mathcal{E}$ has length at most 3): for each equation $x_1 \dots x_n = 1$ with $n \geq 4$, we add some new variables $y_2, \dots, y_{n-2}$ with no rational constraint on y_i ($\mathcal{R}_{y_i} = V$), and we replace the equation $x_1 \dots x_n = 1$ by the equations $y_2 = x_1 x_2$, $y_3 = y_2 x_3$, $\dots$ and $y_{n-2} x_{n-1} x_n = 1$. One can also get rid of equations of length one by forgetting about the corresponding variable.

It is convenient to get rid of equations of length 2. This can be done as follows, using the fact that the set of rational subsets of a virtually free group is a Boolean algebra (Lemma 1.10). If we have an equation $x_1 x_2 = 1$ where x_1 is distinct from $x_2, \bar{x}_2$ as a formal variable, one can substitute every occurrence of x_2 (resp. $\bar{x}_2$) with $\bar{x}_1$ (resp. x_1), and change the rational constraint $x_1 \in \mathcal{R}_{x_1}$ to $x_1 \in \mathcal{R}_{x_1} \cap \mathcal{R}_{x_2}^{-1}$. Of course, one can forget about equations of the form $x\bar{x} = 1$. Each equation of the form $x^2 = 1$ can be replaced by two equations $x^4 = 1$ and $x^6 = 1$, which can be triangulated in the usual way.

The transformation into a triangular system of equations is algorithmic.

2.2 Twisted equations

Consider the general situation of a group V containing a normal subgroup G of finite index. Let $\pi : V \rightarrow Q = V/G$ the quotient map. For each $q \in Q$, choose a lift $\tilde{q} \in V$. Given $x_1, x_2, x_3 \in V$, let $q_i = \pi(x_i)$ and write $x_i = \tilde{q}_i g_i$ for some $g_i \in G$. Then x_1, x_2, x_3 satisfies the equation $x_1 x_2 x_3 = 1$ if and only if $q_1 q_2 q_3 = 1$, (so $\tilde{q}_1 \tilde{q}_2 \tilde{q}_3 \in G$), and $(\tilde{q}_1 \tilde{q}_2 \tilde{q}_3) g_1^{\tilde{q}_2 \tilde{q}_3} g_2^{\tilde{q}_3} g_3 = 1$. This last equation can be viewed as an equation with unknowns g_1, g_2, g_3 , twisted by automorphisms of the form $g \mapsto g^v$ for some $v \in V$.

The group generated by these automorphisms has finite image in $\text{Out}(G)$, but it usually fails to lift to a finite subgroup of $\text{Aut}(G)$.

In Proposition 2.4 below, we prove that when G is a free group, we can embed G in a larger free group $\langle S \rangle$ with free basis S , so that the twisting automorphisms give rise to automorphisms of $\langle S \rangle$ that preserve $S \cup S^{-1}$. In particular, these automorphisms of $\langle S \rangle$ preserve word length, which will be of importance to us. The price to pay is that we have to add rational constraints to our system of equations (even if there were no such constraints originally) to guarantee that the solutions belong to the original free group.

Given a finite set S , and $\langle S \rangle$ the corresponding free group, we denote by $\text{Aut}(S_{\pm})$ the finite group of automorphisms of $\langle S \rangle$ preserving the set $S_{\pm} = S \cup S^{-1}$ (it has order $2^n n!$).

Definition 2.2 (Twisted equations). *A (triangular) system of twisted equations with rational constraints $(\mathcal{E}, \underline{\mathcal{R}})$ in a free group $\langle S \rangle$ consists of an alphabet of variables X , of a finite set $\mathcal{E}$ of equations of the form $((\varphi_1, x_1), (\varphi_2, x_2), (\varphi_3, x_3)) \in (\text{Aut}(S_{\pm}) \times (X \cup \bar{X}))^3$ (representing the equation $\varphi_1(x_1)\varphi_2(x_2)\varphi_3(x_3) = 1$), and a tuple $\underline{\mathcal{R}} = (\mathcal{R}_x)_{x \in X}$ of rational subsets of $\langle S \rangle$.*

A solution of $\mathcal{E}$ is a tuple $\underline{g} = (g_x)_{x \in X} \in \langle S \rangle^X$ with $g_x \in \mathcal{R}_x$ for each $x \in X$, and such that for each $((\varphi_1, x_1), (\varphi_2, x_2), (\varphi_3, x_3)) \in \mathcal{E}$, $\varphi_1(g_{x_1})\varphi_2(g_{x_2})\varphi_3(g_{x_3}) = 1$ where we define $g_{\bar{x}} = g_x^{-1}$ for each $x \in X$.

Remark 2.3. We opted for a restricted meaning of twisting. In general (and as presented in the introduction) a twisted equation is an equation where some automorphisms are involved, but here, the only allowed automorphisms are those that preserve a basis of the given free group. This will be enough for our needs.

The *twisting subgroup* $\Phi \subset \text{Aut}(S_{\pm})$ is the subgroup generated by the twisting morphisms involved in all twisted equations. We say that Φ has no *inversion* if for all $s \in S$ and $\varphi \in \Phi$, $\varphi(s) \neq s^{-1}$. Our construction from a virtually free group will produce only twisting subgroups without inversions. However, if we are given a system of twisted equations where Φ has inversions, one can easily reduce to the case without inversion using barycentric subdivision as follows. Replace $S = \{s_1, \dots, s_n\}$ by $S' = \{u_1, v_1, \dots, u_n, v_n\}$, and embed $\langle S \rangle \rightarrow \langle S' \rangle$ using $s_i \mapsto u_i v_i^{-1}$. If $\Phi(s_i) = s_j$, define $\Phi'(u_i) = u_j$ and $\Phi'(v_i) = v_j$, and if $\Phi(s_i) = s_j^{-1}$, define $\Phi'(u_i) = v_j^{-1}$ and $\Phi'(v_i) = u_j^{-1}$.

Proposition 2.4. *The problem of solving equations with regular constraints in a virtually free group can be reduced to the problem of solving twisted equations in a free group.*

More precisely, let $(\mathcal{E}, \underline{\mathcal{R}})$ be a system of equations with rational constraints in a virtually free group V .

Then there is a free group $\langle S \rangle$, and a finite set $(\mathcal{E}'_1, \underline{\mathcal{R}}'_1) \dots (\mathcal{E}'_k, \underline{\mathcal{R}}'_k)$ of systems of twisted equations (in the sense of Definition 2.2) with rational constraints on $\langle S \rangle$, with the same sets of unknowns X , such that there is a natural bijection between the set of solutions of $\mathcal{E}$ and the (disjoint) union of the sets of solutions of $(\mathcal{E}'_1, \underline{\mathcal{R}}'_1), \dots, (\mathcal{E}'_k, \underline{\mathcal{R}}'_k)$. The twisting subgroup has no inversion and is finite.

Moreover, one can algorithmically compute $\langle S \rangle$ and $(\mathcal{E}'_1, \underline{\mathcal{R}}'_1) \dots (\mathcal{E}'_k, \underline{\mathcal{R}}'_k)$ from a presentation of V , and from $(\mathcal{E}, \underline{\mathcal{R}})$, where all rational subsets are represented by automata.

Remark 2.5. The free group $\langle S \rangle$ does not occur naturally as a subgroup of V . Only a subgroup of $\langle S \rangle$ can be naturally identified as a subgroup of finite index of V . The natural bijection defined in the proof is clearly computable, but we won't need this fact.

Proof. Write V as an extension $F \rightarrow V \xrightarrow{\pi} Q$ where Q is finite and $F = \ker \pi$ is a normal free subgroup of finite index of V . Write V as the fundamental group of a finite graph of finite groups, and let $V \curvearrowright T$ be the dual Bass-Serre tree. By adding a new edge to the graph of groups, we may assume that T contains a point $*$ with trivial stabiliser (T may fail to be minimal).

Consider the graph $\Gamma = T/F$, and still denote by $*$ the image of $*$ in Γ . Since F acts freely, the quotient map $T \rightarrow \Gamma$ is a covering map, and $F = \pi_1(\Gamma, *)$. Note that Q acts on Γ by graph automorphisms (which may fail to fix $*$).

Consider a set S of oriented edges of Γ obtained by choosing an orientation for each edge, and consider the free group $\langle S \rangle$. Each edge path in Γ defines an element of $\langle S \rangle$, and in particular, $F \subset \langle S \rangle$. Clearly, Q acts on $\langle S \rangle$ via automorphisms of $\text{Aut}(S_{\pm})$. We denote by $\varphi_q(x)$ the image of $x \in \langle S \rangle$ under the action of $q \in Q$.

For $g \in V$, let c_g be the edge path of Γ obtained by projection of the segment $[*, g.*] \subset T$. This defines a map $c : V \rightarrow \langle S \rangle$. Since the stabiliser of $*$ is trivial, c is one-to-one. Note that c_g joins $*$ to $\varphi_{\pi(g)}(*)$. The map c is not a morphism but satisfies $c(gh) = c(g) \cdot \varphi_{\pi(g)}(c(h))$ and $c(g^{-1}) = \varphi_{\pi(g)^{-1}}(c(g))$. In particular, the restriction of c to F is a morphism.

Each solution of $\mathcal{E}$ in V maps to a solution of $\mathcal{E}$ in Q . For each solution $\underline{q} \in Q^X$ of $\mathcal{E}$ in Q , the set $\mathcal{R}_x \cap \pi^{-1}(q_x)$ is a rational subset by Lemma 1.10. Let $\mathcal{S}_{\underline{q}} \subset V^X$ be the set of solutions of $\mathcal{E}$ with rational constraints $x \in \mathcal{R}_x \cap \pi^{-1}(q_x)$. The set of solutions of $(\mathcal{E}, \underline{\mathcal{R}})$ is the disjoint union of the sets $\mathcal{S}_{\underline{q}}$.

Each equation in $\mathcal{E}$ is equivalent to an equation of the form $x_1 x_2 x_3 = 1$ or $x_1 x_2 \bar{x}_3 = 1$ where x_1, x_2, x_3 lie in X (not $\bar{X}$); we assume that each equation is of this form. Given $\underline{g} \in \mathcal{S}_{\underline{q}}$, the equation $g_{x_1} g_{x_2} g_{x_3} = 1$ implies $c(g_{x_1}) \cdot \varphi_{q_{x_1}}(c(g_{x_2})) \cdot \varphi_{q_{x_1} q_{x_2}}(c(g_{x_3})) = 1$. Similarly, the equation $g_{x_1} g_{x_2} g_{x_3}^{-1} = 1$ implies $c(g_{x_1}) \cdot \varphi_{q_{x_1}}(c(g_{x_2})) \cdot c(g_{x_3})^{-1} = 1$.

Let $\mathcal{E}'_{\underline{q}}$ be the set of equations in $\langle S \rangle$ obtained by replacing in $\mathcal{E}$ each equation $x_1 x_2 x_3 = 1$ (resp. $x_1 x_2 \bar{x}_3 = 1$) by the twisted equation $x_1 \cdot \varphi_{q_{x_1}}(x_2) \cdot \varphi_{q_{x_1} q_{x_2}}(x_3) = 1$ (resp. $x_1 \cdot \varphi_{q_{x_1}}(x_2) \cdot \bar{x}_3 = 1$). Thus c maps $\mathcal{S}_{\underline{q}}$ into the set $\mathcal{S}'_{\underline{q}}$ of solutions of $(\mathcal{E}'_{\underline{q}}, \underline{\mathcal{R}}')$ where $\mathcal{R}'_{\underline{q}, x} = c(\mathcal{R}_x \cap \pi^{-1}(q_x)) \subset \langle S \rangle$. Let's check that $\mathcal{R}'_{\underline{q}, x}$ is a rational subset of $\langle S \rangle$. If $q_x = 1$, this follows from the fact that $c|_F$ is a morphism. Otherwise, consider $\tilde{q}_x \in \pi^{-1}(\{q_x\})$, and consider the rational subset $\mathcal{R}'' = \tilde{q}_x^{-1}(\mathcal{R}_x \cap \pi^{-1}(q_x)) \subset F$. Since $c|_F$ is a morphism, $c(\mathcal{R}'')$ is rational and so is $\mathcal{R}'_{\underline{q}, x} = c(\tilde{q}_x) \cdot \varphi_{q_x}(c(\mathcal{R}''))$.

We claim that c maps $\mathcal{S}_{\underline{q}}$ onto $\mathcal{S}'_{\underline{q}}$. Indeed, consider a solution $\underline{a} \in \mathcal{S}'_{\underline{q}}$. For each $x \in X$, since $a_x \in \mathcal{R}'_{\underline{q}, x}$, a_x corresponds to a path joining $*$ to $q_x.*$. This path lifts to a path in the tree T joining $*$ to some $g_x.*$, where g_x is uniquely determined because $*$ has trivial stabiliser. By definition, $c(g_x) = a_x$. Since c realises a bijection between $\mathcal{R}_x \cap \pi^{-1}(q_x)$ and its image, the constraint $g_x \in \mathcal{R}_x \cap \pi^{-1}(q_x)$ is satisfied. Since for each equation $x_1 x_2 x_3 \in \mathcal{E}$, $\underline{a}$ satisfies the corresponding equation in $\mathcal{E}'_{\underline{q}}$, namely $a_{x_1} \cdot \varphi_{q_{x_1}}(a_{x_2}) \cdot \varphi_{q_{x_1} q_{x_2}}(a_{x_3}) = 1$, we get that $c(g_{x_1} g_{x_2} g_{x_3}) = 1$. By injectivity of c , $g_{x_1} g_{x_2} g_{x_3} = 1$. A similar argument applies to equations of the form $x_1 x_2 \bar{x}_3 \in \mathcal{E}$.

This proves that c induces a bijection between the set of solutions of $(\mathcal{E}, \underline{\mathcal{R}})$ and the (disjoint) union of the sets of solutions of $(\mathcal{E}'_{\underline{q}}, \underline{\mathcal{R}}'_{\underline{q}})$ as $\underline{q}$ ranges among solutions of $\mathcal{E}$ in Q^X .

Let's prove the computability of the new system of equations. Starting with a presentation of a virtually free group V , one can effectively find a presentation of V as a graph of finite groups Λ . Indeed, enumerating all presentations of V , one will find one which is visibly a presentation of the correct form. More precisely, if A is a finite group, its finite presentation $\langle A | \text{table}(A) \rangle$ consisting of its multiplication table is a presentation from which finiteness of A is obvious. Consider an amalgam of two finite groups $G = A *_C B$, given by two monomorphisms $j_A : C \rightarrow A$, $j_B : C \rightarrow B$. Then G has a presentation of the form $\langle A \cup B \cup C | \text{table}(A), \text{table}(B), \text{table}(C), \{c = j_A(c) = j_B(c)\}_{c \in C} \rangle$ from which one can obviously read that j_A, j_B are injective morphisms, and that G is an amalgam of finite groups. Similarly, given a graph of finite groups Γ and $\tau \subset \Gamma$ a maximal tree, one produces a presentation of $\pi_1(\Gamma, \tau)$ from multiplication tables of vertex and edge groups from which one can read that the corresponding group is the fundamental group of a graph of finite groups.

One can also find a normal free subgroup F of finite index: enumerate all morphisms to finite groups $\rho : V \rightarrow Q$, and check whether for each vertex group Λ_v , $\rho|_{\Lambda_v}$ is injective; one will eventually find such a ρ , and one can take $F = \ker \rho$. The graph $\Gamma = T/F$ is constructed from Λ as the covering with deck group Q : for each edge or

vertex x of Λ , we put a copy of this edge or vertex for each element of $Q/\rho(\Lambda_x)$, and the incidence relation is the natural one. Thus $\langle S \rangle$ is computable together with its natural basis and the action of Q . The new systems of equations $\mathcal{E}'_q$ are now explicit.

There remains to compute the rational constraints $\mathcal{R}'_{q,x} = c(\mathcal{R}_x \cap \pi^{-1}(q_x))$. We saw that $\mathcal{R}'_{q,x} = c(\tilde{q}_x) \cdot \varphi_{q_x}(c(\mathcal{R}''))$ with $\mathcal{R}'' = \tilde{q}_x^{-1}(\mathcal{R}_x \cap \pi^{-1}(q_x))$. An automaton representing $\mathcal{R}''$ can be computed using Lemma 1.10. By Lemma 1.4, since $c|_F$ is a morphism, we get an automaton representing $c(\mathcal{R}'')$, and thus $\mathcal{R}'_{q,x}$. $\square$

3 Band complexes

We fix a finite set S , the free group $G = \langle S \rangle$, and the corresponding free monoid $S_\pm^*$ with involution $x \rightarrow \bar{x}$. Let $\text{Aut}(S_\pm)$ be the corresponding finite group of automorphisms of G preserving $S \cup S^{-1}$, which we identify with the group of automorphisms of $S_\pm^*$ commuting with the involution.

3.1 Band complexes

A *band complex* Σ consists of a *domain* D together with a finite set of bands. The domain D is a disjoint union of non-degenerate segments. We say that a segment is non-degenerate when it is non-empty and not reduced to a point. A *band* B consists of a rectangle $[a, b] \times [0, 1]$, together with two injective continuous *attaching* maps $f_{B,0} : [a, b] \times \{0\} \rightarrow D$ and $f_{B,1} : [a, b] \times \{1\} \rightarrow D$ and a *twisting* automorphism $\varphi_B \in \text{Aut}(S_\pm)$. The segments $J_{B,\varepsilon} = f_{B,\varepsilon}([a, b] \times \{\varepsilon\})$ for $\varepsilon = 0, 1$ are called *the bases* of the band B .

Remark. Since bands carry twisting automorphisms, a band complex might rather be called a *twisted* band complex. We won't use this terminology for shortness's sake.

We denote by $\Phi \subset \text{Aut}(S_\pm)$ the group generated by all twisting morphisms φ_B , and we will assume that Φ has no inversion, i. e. that $\varphi(s) \neq \bar{s}$ for all $\varphi \in \Phi$ and $s \in S_\pm$.

The set of *vertices* of Σ is the subset of D consisting of the endpoints of D together with the endpoints of the bases of the bands. We identify Σ with the CW-complex obtained by gluing the bands on D using the maps $f_{B,\varepsilon}$.

The precise value of the attaching maps are not important: a band complex is really determined by the ordering of vertices in each component of D , and for each band, the 4-tuple $(f_{B,0}(a), f_{B,0}(b), f_{B,1}(a), f_{B,1}(b))$ together with the twisting automorphism φ_B .

3.2 Solution of a band complex

An *elementary segment* of Σ is a segment of D joining two adjacent vertices. We denote by $E(\Sigma)$ the set of oriented elementary segments of Σ . We denote by $e \mapsto \bar{e}$ the orientation reversing involution on $E(\Sigma)$.

A *labelling* σ of Σ is a labelling of each $e \in E(\Sigma)$ by a non-empty word $\sigma_e \in S_\pm^+$, so that $\sigma_{\bar{e}} = \overline{\sigma_e}$ (we use the standard notation $S_\pm^+ = S_\pm \cdot S_\pm^* = S_\pm^* \setminus \{1\}$). Say that a segment $I \subset D$ is *adapted* to Σ if it is a union of elementary segments. For each oriented adapted segment $I = e_1 \dots e_n \subset D$ written as a concatenation of elementary segments, we define $\sigma_I = \sigma_{e_1} \dots \sigma_{e_n}$. Alternatively, we often view a labelling of Σ as

a subdivision of D , together with a labelling of the subdivided edges by letters in $S \cup \overline{S}$.

A *solution* σ of Σ is a labelling such that for each band B with twisting automorphism φ_B and with bases J_0, J_1 oriented so that the attaching maps preserve the orientation, one has $\sigma_{J_1} = \varphi_B(\sigma_{J_0})$. We say that the solution σ is *reduced* if for every non-degenerate adapted segment $I \subset D$, σ_I is a non-trivial reduced word.

The *length* $|\sigma|$ of a solution σ is the sum of lengths of words $|\sigma_{D_1}| + \dots + |\sigma_{D_k}|$ where $D_1, \dots, D_k$ are the connected components of D (with a choice of orientation).

3.3 Rational constraints

A set of *rational constraints* on Σ is a family of regular languages $\underline{\mathcal{R}} = (\mathcal{R}_e)_{e \in E(\Sigma)}$ of $S_{\pm}^+$, indexed by oriented elementary segments of Σ , and such that $\mathcal{R}_{\bar{e}} = \overline{\mathcal{R}_e}$. A *solution* σ of Σ , satisfying the rational constraint $\underline{\mathcal{R}}$ is a solution of Σ such that for each elementary segment $e \in E(\Sigma)$, $\sigma_e \in \mathcal{R}_e$.

All the rational constraints we will use later will be in some *standard form* as defined in Section 3.5.

3.4 From twisted equations to band complexes

Proposition 3.1. *Let $(\mathcal{E}, \underline{\mathcal{R}})$ be a system of twisted equations with rational constraints on a free group $\langle S \rangle$. Then one can effectively compute a finite set of band complexes with rational constraints $\Sigma_1, \dots, \Sigma_p$ and a bijection between the set of solutions of $(\mathcal{E}, \underline{\mathcal{R}})$ and the disjoint union of the (reduced) solutions of $\Sigma_1, \dots, \Sigma_p$.*

Moreover, every solution of Σ_i is reduced.

Proof. Say that a tuple $g \in G^X$ is *singular* if $g_x = 1$ for some $x \in X$. Denote by $\text{Sol}(\mathcal{E}, \underline{\mathcal{R}})$ (resp. $\text{Sol}^*(\mathcal{E}, \underline{\mathcal{R}})$) the set of solutions (resp. of non-singular solutions) of $(\mathcal{E}, \underline{\mathcal{R}})$. Clearly, $\text{Sol}(\mathcal{E}, \underline{\mathcal{R}})$ is in bijection with the disjoint union of $2^{|X|}$ sets of the form $\text{Sol}^*(\mathcal{E}_{X_0}, \underline{\mathcal{R}})$ where for $X_0 \subset X$, $\mathcal{E}_{X_0}$ is the system of twisted equations whose set of unknowns is $X \setminus X_0$, and obtained from $\mathcal{E}$ by replacing each occurrence of the variable $x \in X_0$ by 1.

Some of the obtained equations may involve less than three variables. However, if some equation involves just one variable, it is of the form $\varphi(x) = 1$, so $\mathcal{E}$ has no non-singular solution and we may forget about this X_0 . We view the equations involving two variables as twisted equalities of the form $\varphi_1(x_1) = \varphi_2(x_2)$.

Remark 3.2. We could get rid of twisted equalities involving two distinct formal variables by substitution, and of twisted equalities of the form $\varphi(x) = x$ by adding the rational constraint saying that $x \in \text{Fix}\varphi$, but this does not allow to get rid of equations of the form $\varphi(\bar{x}) = x$ since the set of fixed points of $x \mapsto \varphi(\bar{x})$ is not a rational language in general, and we don't want to add new singular variables.

We now need to build some band complexes encoding the set of non-singular solutions of a given system of twisted equations $(\mathcal{E}, \underline{\mathcal{R}})$ (including twisted equalities).

For each unknown $x \in X$, we consider an oriented segment D_x (whose labelling will correspond to the value g_x of the unknown x). For $x \in \overline{X}$, we will denote by $D_{\bar{x}}$ the same segment as D_x , but with the opposite orientation. Then for each twisted equation ε of the form $\varphi_1(x_1)\varphi_2(x_2)\varphi_3(x_3) = 1$ (resp. $\varphi_1(x_1) = \varphi_2(x_2)$), with $x_i \in X \cup \overline{X}$, we add three (resp. two) oriented segments $D_{\varepsilon, i}$ corresponding to $\varphi_i(x_i)$. We define the domain D of our band complex as $D = (\bigsqcup_{x \in X} D_x) \sqcup (\bigsqcup_{(\varepsilon, i)} D_{\varepsilon, i})$.

For each segment $D_{\varepsilon,i}$ corresponding to $\varphi_i(x_i)$, we add a band whose bases are $J_0 = D_{x_i}$ and $J_1 = D_{\varepsilon,i}$, preserving the orientation, and whose twisting morphism is φ_i .

For each twisted equality ε of the form $\varphi_1(x_1) = \varphi_2(x_2)$, we add a band whose bases are $J_0 = D_{\varepsilon,1}$ and $J_1 = D_{\varepsilon,2}$, preserving the orientation, and with trivial twisting morphism.

For any non-singular solution $\underline{g}$ of the twisted equation $\varepsilon \in \mathcal{E}$ corresponding to $\varphi_1(x_1)\varphi_2(x_2)\varphi_3(x_3) = 1$, one can consider its cancellation tripod τ : this is the convex hull of $a_1 = 1, a_2 = \varphi_1(g_{x_1}), a_3 = \varphi_1(g_{x_1})\varphi_2(g_{x_2})$ in the Cayley graph of $\langle S \rangle$, so that $[a_i, a_{i+1}]$ is labelled by $\varphi_i(g_{x_i})$ (where we view i as an integer mod 3). Let c be the centre of τ , i. e. the intersection of the three segments $[a_i, a_{i+1}]$. There are four possible types of combinatorics of cancellation for each triangular equation ε : $c \notin \{a_1, a_2, a_3\}$ (in which cases τ is not a segment), or $c = a_1, c = a_2, c = a_3$. The four types are mutually exclusive because $\underline{g}$ is non-singular.




Figure 2: The bands when the cancellation tripod is non-degenerate

To each choice of combinatorics of cancellation for this twisted equation, we associate a set of bands to be added to our band complex. If the cancellation tripod is not a segment, for each $i = 1, \dots, 3$, add a vertex $c_{\varepsilon,i}$ at the midpoint of $D_{\varepsilon,i}$ (Figure 2). Then add a band reversing the orientation, whose bases are the right half-segment of $D_{\varepsilon,i}$ and the left half-segment of $D_{\varepsilon,i+1}$ (right and left having a meaning relative to the orientation of those segments), and whose twisting automorphism is the identity.

If the cancellation tripod is such that $c = a_i$ for some $i \in \{1, 2, 3\}$, the product $\varphi_{i-1}(g_{x_{i-1}})\varphi_i(g_{x_i})$ is reduced, i being thought modulo 3 (Figure 3). We add one vertex at the midpoint of $D_{\varepsilon,i+1}$, and two bands reversing the orientation, whose bases are $J_0 = D_{\varepsilon,i}$ (resp. $J_0 = D_{\varepsilon,i-1}$) and J_1 the initial (resp. final) half-segment of $D_{\varepsilon,i+1}$, and whose twisting automorphism is the identity.

Let t be the number of triangular equations of $\mathcal{E}$. For each choice κ out of the 4^t possible combinatorics of cancellation, we obtain a band complex Σ_κ .

Finally, for each rational subset $\mathcal{R}_x \subset \langle S \rangle$, we consider $\tilde{\mathcal{R}}_x \subset S_\pm^*$ the set of reduced words representing elements of $\mathcal{R}_x$. This is a rational language by Lemma 1.6. We add to Σ_κ the rational constraint $\tilde{\mathcal{R}}_x$ on D_x for each $x \in X$, and we don't put any constraint on the other elementary segments of Σ (i. e. we set $\mathcal{R}_e = S_\pm^*$).




Figure 3: The bands when the cancellation tripod is flat

Since $\tilde{\mathcal{R}}_x$ is a language of reduced words, any solution of Σ_κ is reduced.

We claim that the set of reduced solutions of Σ_κ is in one-to-one correspondence with the subset $\mathcal{S}_\kappa$ of $\text{Sol}^*(\mathcal{E}, \underline{\mathcal{R}})$ whose combinatorics of cancellation correspond to κ .

Indeed, a non-singular solution $\underline{g} \in \mathcal{S}_\kappa$ defines a labelling σ of Σ_κ as follows: D_x is labelled by the reduced word $\sigma_x \in S_\pm^*$ representing g_x , each $D_{\varepsilon,i}$ representing $\varphi_i(x_i)$ is labelled by the reduced word representing $\varphi_i(g_{x_i})$, and if the midpoint of $D_{\varepsilon,i}$ is a vertex of Σ_κ , the labelling of the two half segments of $D_{\varepsilon,i}$ should be such that the position of this vertex corresponds to the centre of the cancellation tripod. The labelling σ thus constructed is a clearly reduced solution of Σ_κ .

Conversely, if σ is a reduced solution of Σ_κ , the image g_x in $\langle S \rangle$ of the word σ_{D_x} clearly defines a non-singular solution of $(\mathcal{E}, \underline{\mathcal{R}})$. Moreover, for each $\varepsilon \in \mathcal{E}_T$, the label of $D_{\varepsilon,i}$ defines a geodesic segment, and the midpoint of the three segments $D_{\varepsilon,i}$ corresponds to the centre of the cancellation tripod for ε . Thus, the combinatorics of cancellation agree with κ , and $g_x \in \mathcal{S}_\kappa$.

The construction of the band complexes Σ_κ is clearly effective, so Proposition 3.1 follows. $\square$

3.5 Standard forms of rational constraints

It will be convenient to represent all the needed rational languages in a fixed finite monoid. This will allow to treat uniformly all rational languages appearing under various transformations of the band complexes.

Lemma 3.3. *Let $\mathcal{R}_1, \dots, \mathcal{R}_k$ be a finite set of rational languages in $S_\pm^+$.*

There exists a finite monoid $\mathcal{M}$ with an involution $m \mapsto \overline{m}$ and with an action of $\text{Aut}(S_\pm)$, and an $\text{Aut}(S_\pm)$ -equivariant morphism $\rho : S_\pm^+ \rightarrow \mathcal{M}$ commuting with the involutions, such that each $\mathcal{R}_i$ is the preimage of a finite subset of $\mathcal{M}$.

Moreover, all this data is algorithmically computable from automata representing $\mathcal{R}_i$.

When $\mathcal{R}$ is the preimage under ρ of some subset of $\mathcal{M}$ as above, we say that $\mathcal{R}$ is represented by ρ .

Proof. We first consider the case of a single regular language. Let $\mathcal{R} \subset S_{\pm}^+$ be a rational language. Let A be an automaton representing L with vertex set $v_1, \dots, v_n$. For each $s \in S \cup \bar{S}$, consider the $n \times n$ Boolean incidence matrix M_s of the subgraph of A whose edges are those labelled by s .

Let $\mathcal{M}_0$ be the semigroup of $n \times n$ -Boolean matrices, and $\rho_0 : S_{\pm}^+ \rightarrow \mathcal{M}_0$ the morphism sending s to M_s . Let v_S (resp. v_A) be the Boolean vector representing *Start* (resp. *Accept*). Then $\mathcal{R}$ is the preimage under ρ of the set $A_0 \subset \mathcal{M}_0$ of matrices M such that ${}^t v_S M v_A = 1$.

Now let $\bar{\rho}_0 : S_{\pm}^+ \rightarrow \mathcal{M}_0$ be the morphism sending s to ${}^t M_{\bar{s}}$. Note that $\bar{\rho}_0(w) = {}^t \rho_0(\bar{w})$ for all $w \in S_{\pm}^+$. Consider the finite monoid $\mathcal{M} = \mathcal{M}_0 \times \mathcal{M}_0$ endowed with the involution $(M_1, M_2) \mapsto \overline{(M_1, M_2)} \stackrel{\text{def}}{=} ({}^t M_2, {}^t M_1)$, and consider $\rho : S_{\pm}^+ \rightarrow \mathcal{M}$ sending s to $(\rho_0(s), \bar{\rho}_0(s))$. By the remark above, ρ commutes with the involutions of $S_{\pm}^+$ and $\mathcal{M}$, and $\mathcal{R} = \rho^{-1}(A_0 \times \mathcal{M}_0)$.

Now given a finite set of languages $\mathcal{R}_1, \dots, \mathcal{R}_k$, consider finite monoids with involutions $\mathcal{M}_1, \dots, \mathcal{M}_k$ and $\rho_i : S_{\pm}^+ \rightarrow \mathcal{M}_i$ commuting with the involutions such that $\mathcal{R}_i = \rho_i^{-1}(A_i)$ for some $A_i \subset \mathcal{M}_i$. Consider the finite monoid $\mathcal{M} = \mathcal{M}_1 \times \dots \times \mathcal{M}_k$ with the diagonal involution, and $\rho = \rho_1 \times \dots \times \rho_k$. Then for $A'_i = \mathcal{M}_1 \times \dots \times \mathcal{M}_{i-1} \times A_i \times \mathcal{M}_{i+1} \times \dots \times \mathcal{M}_k$, $\mathcal{R}_i = \rho^{-1}(A'_i)$.

Finally, we need to put the $\text{Aut}(S_{\pm})$ -action in the picture. For each $\alpha \in \text{Aut}(S_{\pm})$, let $\mathcal{M}_{\alpha}$ be a copy of $\mathcal{M}$, and consider $\rho_{\alpha} = \rho \circ \alpha : S_{\pm}^+ \rightarrow \mathcal{M}_{\alpha}$. Consider the product $\tilde{\mathcal{M}} = \prod_{\alpha \in \text{Aut}(S_{\pm})} \mathcal{M}_{\alpha}$ endowed with the action of $\text{Aut}(S_{\pm})$ by permutation of factors, and $\tilde{\rho} = \prod_{\alpha \in \text{Aut}(S_{\pm})} \rho_{\alpha}$. Then $\tilde{\rho} : S_{\pm}^+ \rightarrow \tilde{\mathcal{M}}$ satisfies the lemma. $\square$

By Lemma 3.3, one can represent all the rational constraints $\mathcal{R}_e$ of a given band complex by a single morphism $\rho : S_{\pm}^+ \rightarrow \mathcal{M}$. It will be useful to assume that ρ also represents $\text{Fix}\varphi \subset S_{\pm}^+$ for all $\varphi \in \Phi$ (this is a regular language as it is the submonoid generated by Φ -invariant letters in $S_{\pm}$). By a finite disjunction of cases, we need only to consider rational constraints of the form $\mathcal{R}_e = \rho^{-1}(\{m_e\})$.

Definition 3.4 (Standard form for rational constraints). *A set of rational constraints in standard form on a band complex Σ consists of*

1. *a finite monoid $\mathcal{M}$ with an involution and an action of $\text{Aut}(S_{\pm})$,*
2. *an epimorphism $\rho : S_{\pm}^+ \rightarrow \mathcal{M}$ commuting with the involutions and $\text{Aut}(S_{\pm})$ -equivariant, such that for all $\varphi \in \text{Aut}(S_{\pm})$, the rational language $\text{Fix}\varphi \subset S_{\pm}^+$ is represented by ρ ,*
3. *a collection $\underline{m} = (m_e)_{e \in E(\Sigma)} \in \mathcal{M}^{E(\Sigma)}$ such that $m_{\bar{e}} = \overline{m_e}$.*

The tuple $\underline{m} = (m_e)$ defines a tuple of rational constraints by $\mathcal{R}_e = \rho^{-1}(\{m_e\})$. By definition, a solution of $(\Sigma, \underline{m})$ is a solution of Σ with the corresponding rational constraints. Using Lemma 3.3, we immediately get the following lemma. The fact that ρ is onto can be obtained by changing $\mathcal{M}$ to $\rho(S_{\pm}^+)$.

Lemma 3.5. *Given a band complex Σ with rational constraints $\underline{\mathcal{R}}$, one can compute a finite monoid $\mathcal{M}$, a morphism $\rho : S_{\pm}^+ \rightarrow \mathcal{M}$, and some tuples $\underline{m}_1, \dots, \underline{m}_p \in \mathcal{M}^{E(\Sigma)}$ defining rational constraints in standard form on Σ such that the set of solutions of $(\Sigma, \underline{\mathcal{R}})$ is the disjoint union of the sets of solutions of $(\Sigma, \underline{m}_1), \dots, (\Sigma, \underline{m}_p)$.*

From now on, all band complexes are endowed with rational constraints in standard form.

4 Bulitko's Lemma: bounding the exponent of periodicity

The goal of this section is a version of Bulitko's lemma [Bul70] adapted to twisted equations, which we state in terms of band complexes. The *exponent of periodicity* of a solution σ of a band complex Σ is the largest integer s such that some subsegment of the domain D of Σ is labelled by some word of the form p^s , with p a non-trivial word. Recall that the length of a solution σ of a band complex is the number of letters in $S_{\pm}$ involved in the labelling.

Proposition 4.1 ([Bul70]). *Let Σ be a band complex with rational constraints $\underline{m}$ in standard form.*

There exists a computable number $B(\Sigma, \underline{m})$ such that any shortest solution of Σ has an exponent of periodicity at most $B(\Sigma, \underline{m})$.

We will use and prove the proposition only for a band complex arising from a system of equations as in Section 3.4. More precisely, we assume that every component D_0 of the domain of the band complex contains at most one vertex in its interior, and at most two bases of bands distinct from D_0 (see Figure 2 and 3). The general statement can be easily deduced from this case.

4.1 Normal decomposition with respect to powers

Consider a solution σ of the band complex Σ . We view σ as a subdivision of D together with a labelling of the edges by elements of $S_{\pm}$. Each interval $A \subset D$ we consider is a union of such edges. Its length $|A|$ is the number of edges it contains, which coincides with the length of the word σ_A labelled by A .

We fix some $q \in \mathbb{N} \setminus \{0\}$. Given some interval $A \subset D$, we are going to define a natural set of *special segments* encoding its large enough subwords which are q -periodic. Say that an interval $A \subset D$ of length $n > q$ is *q -periodic* if $\sigma_A = a_1 \dots a_n$ with $a_i \in S_{\pm}$ and for all $i \leq n - q$, $a_i = a_{i+q}$.

For each maximal q -periodic segment $P \subset A$ of length at least $2q$, we say that its central subsegment $Q \subset P$ of length $|P| - 2q$ is a *special segment* of A . A special segment is reduced to a vertex when $|P| = 2q$. Although maximal q -periodic segments may overlap, the following claim holds.

Claim 4.2. *Two distinct q -special segments of A are disjoint, and separated by a segment of length at least $q + 1$.*

Proof. Let $Q, Q' \subset A$ be two special segments, and $P \supset Q$ and $P' \supset Q'$ the corresponding maximal q -periodic segments. If the claim fails, then $P \cap P'$ has length at least q . Thus, two edges of $Q \cup Q'$ at distance q of each other either both lie in P or in P' , so $P \cup P'$ is q -periodic. $\square$

For any word p of length q , and $r = n/q \in \frac{1}{q}\mathbb{N}$, it will be convenient to denote by p^r the prefix of length n of a sufficiently large power of p . If $r \in \mathbb{N}$, then p^r coincides with the usual r -th power of p . For each special segment Q we denote by $r(Q) = |Q|/q \in \frac{1}{q}\mathbb{N}$ the exponent and by $p(Q)$ the period so that Q is labelled by $p(Q)^{r(Q)}$ (if $r(Q) < 1$, $p(Q)$ should be defined using the maximal q -periodic subsegment P containing Q).

The *normal decomposition* of $A \subset D$ (with respect to q and to the solution σ) is the decomposition $A = U_0.Q_1.U_1 \dots Q_n.U_n$ defined by its special segments $Q_1, \dots, Q_n$. Note that if the decomposition is non-trivial (i. e. if $n \geq 1$), then $|U_0| \geq q$ since the suffix of length q of U_0 is $p(Q_1)$. If $\varphi \in \text{Aut}(S_\pm)$, and if $\sigma_{A'} = \varphi(\sigma_A)$ (resp. if $\sigma_{A'} = \sigma_A^{-1}$), then the normal decompositions match (up to changing the orientation).

4.2 Special segments and concatenation

Assume that A, A', A'' are oriented intervals of D such that $A = A'.A''$. Any special segment Q' of A' (or A'') is clearly contained in a special segment Q of A . If $Q = Q'$, we say that Q' is *regular*, and *exceptional* otherwise. We say that a special segment Q of A is *regular* if it is contained in A' or A'' and is a special segment in it.

Let $A' = U'_0.Q'_1.U'_1 \dots Q'_{n'}.U'_{n'}$ and $A'' = U''_0.Q''_1.U''_1 \dots Q''_{n''}.U''_{n''}$ be the normal decomposition of A' and A'' . If Q' (resp. Q'') is an exceptional segment of A' (resp. A''), then $Q' = Q'_{n'}$ (resp. $Q'' = Q''_1$) and by maximality of special segments, $|U'_{n'}| = q$ (resp. $|U''_0| = q$).

Let Q be an exceptional special segment of A . We distinguish several cases.

1. If Q contains two special segments of A' or A'' , then $Q = Q'_{n'}.U'_{n'}.U''_0.Q''_0$ so $r(Q) = r(Q'_{n'}) + r(Q''_0) + 2$.
2. Otherwise, if Q contains exactly one special segment of A or A' , and if this special segment lies in A' , then $Q \supset Q'_{n'}$ and $|Q \cap A''| < q$ since otherwise, $Q \cap A''$ would contain a special segment of A'' . Thus, $r(Q) < r(Q'_{n'}) + 2$.
3. Symmetrically, if Q contains exactly one special segment, lying in A'' , then $Q \supset Q''_0$ and $r(Q) < r(Q''_0) + 2$.
4. If Q contains no special segment of A' or A'' , then $|Q \cap A'| < q$ and $|Q \cap A''| < q$ so $r(Q) < 2$.

There can be at most 2 exceptional segments in A because two special segments are at least at distance $q + 1$ apart. It may happen that there are two exceptional segments: for $q = 5$, consider the word $p = ababa$, A' labelled by $cpaba$ and A'' by $babapc$ so that A' and A'' contain no special segment; then $A'.A''$ is labelled by $cp^2bapc = cpabp^2c$ and has two special segments of length 0.

4.3 Proof of Bulitko's Lemma

We are now ready for the proof of Proposition 4.1. Once normal forms have been adapted to twisted equations, the proof is identical to the one given in [DGH05]. We give it here for the reader's convenience.

Proof. Start with a shortest solution σ of Σ with presumably high exponent of periodicity, i. e. containing high powers of a word of length q . For each special segment Q , recall that $p(Q)$ denotes the word labelled by Q , and $r(Q) = |Q|/q \in \frac{1}{q}\mathbb{N}$. We also define $t(Q) \in \mathbb{N}$ and $u(Q) \in \{0, \frac{1}{q}, \dots, \frac{q-1}{q}\}$ as the integer and fractional part of $r(Q)$ so that $r(Q) = t(Q) + u(Q)$.

The structure of the proof is as follows. We view the $t(Q)$'s as variables satisfying a system of integer equations Λ . Each solution of Λ provides a new solution of Σ ,

and the fact that σ is shortest says that this solution of Λ is minimal. The system of equations happens to be equivalent to a system of equations whose number of equations, variables, and coefficients are bounded independently of σ , thus giving a bound on the size of a minimal solution of Λ , and thus on the exponents of special segments.

Since we assume that our band complex Σ comes from the construction in section 3.4, each component D_0 of its domain D contains at most one vertex in its interior, and at most two bases of bands distinct from D_0 . We first forget about rational constraints.

Let $\mathcal{B}$ be the set of bases of bands of Σ , with a chosen orientation. For each $A \in \mathcal{B}$, let $A = U_{A,0} \cdot Q_{A,1} \cdot U_{A,1} \dots Q_{A,n_A} \cdot U_{A,n_A}$ be its normal decomposition. Given a set of values $(\tau_{A,i}) \in \mathbb{N}^{n_A}$, we define a new word by changing the integer part the exponents of the special segments of A :

$$w_A(\tau_{A,1}, \dots, \tau_{A,n_A}) = \sigma_{U_{A,0}} \cdot (p(Q_{A,1})^{\tau_{A,1}+u(Q_{A,1})}) \cdot \sigma_{U_{A,1}} \dots (p(Q_{A,n_A})^{\tau_{A,n_A}+u(Q_{A,n_A})}) \cdot \sigma_{U_{A,n_A}}$$

By definition, $w_A(t(Q_{A,1}), \dots, t(Q_{A,n_A})) = \sigma_A$.

We now define a system of equations Λ with unknowns $(\tau_{A,k}) \in \bigoplus_{A \in \mathcal{B}} \mathbb{N}^{n_A}$. By construction, $\tau_{A,k} = t(Q_{A,k})$ will be a solution of Λ . For each band with bases A_0, A_1 with twisting morphism φ , we have $\sigma_{A_1} = \varphi(\sigma_{A_0})$, $n_{A_1} = n_{A_0}$, and we add the equations $\tau_{A_1,k} = \tau_{A_0,k}$ for $k \in \{1, \dots, n_{A_1}\}$. For each connected component A of D containing a vertex distinct from its endpoints, decomposing A into $A' \cdot A''$, we add to Λ an equation $\tau_{A',k} = \tau_{A,k}$ for each regular special segment of A' , and an equation $\tau_{A'',n_{A''}-k} = \tau_{A,n_A-k}$ for each regular special segment of A'' .

If Q is a exceptional segment of A containing two exceptional segments $Q' \subset A'$ and $Q'' \subset A''$, then $r(Q) = r(Q') + r(Q'') + 2$, so either $t(Q) = t(Q') + t(Q'') + 2$ or $t(Q) = t(Q') + t(Q'') + 3$; we add to Λ the corresponding equation $\tau_{A,i} = \tau_{A',n_{A'}} + \tau_{A'',1} + 2$ or $\tau_{A,n_A} = \tau_{A',n_{A'}} + \tau_{A'',1} + 3$.

If Q contains exactly one special segment Q' , lying in A' , then $r(Q) < r(Q') + 2$, so $t(Q) = t(Q') + \varepsilon$ for some $\varepsilon \in \{0, 1, 2\}$; we add to Λ the equation $\tau_{A,k} = \tau_{A',n_{A'}} + \varepsilon$ where k is the index corresponding to Q in A . We proceed symmetrically if Q contains exactly one special segment in A'' .

If Q contains no special segment of A' or A'' , then $r(Q) < 2$ so $t(Q) = \varepsilon$ for some $\varepsilon \in \{0, 1\}$, and we add to Λ the equation $\tau_{A,k} = \varepsilon$ where k is the index corresponding to Q in A .

This defines Λ . This system has a solution, namely $\tau_{A,k} = t(Q_{A,k})$. Moreover, if $(\tau_{A,k})$ is any solution of Λ , then $w_A(\tau_{A,1}, \dots, \tau_{A,n_A})$ defines a solution of Σ . In particular, if $t(Q_{A,k})$ is not a minimal solution of Λ , meaning that there exists a solution of Λ with $\tau_{A,k} \leq t(Q_{A,k})$ for all A, k , and some inequality is strict, then σ is not the shortest solution.

The number of equations and of unknowns of Λ is not bounded a priori. However, there are at most $2\#\pi_0(D)$ equations which are not equalities of the type $\tau_{A,k} = \tau_{A',k'}$. By substitution, one can get rid of these equalities, and we are left with a system Λ' of at most $2\#\pi_0(D)$ equations, with coefficients in $\{0, \dots, 3\}$, and with $l \leq 6\#\pi_0(D)$ unknowns. Note that the bounds are independent of σ and of q .

Knowing only the band complex Σ (and not the solution σ), one can list all possible systems Λ' , decide which of them have a solution in $\mathbb{N}^l$, list all minimal solutions of each of them (see for instance [Sch86, Th 17.1]) and compute the maximal value

M of any $\tau_{A,k}$ in any such minimal solution. The value M is then an upper bound on the exponent of periodicity of a shortest solution σ of Σ .

In presence of rational constraints, represented by $\rho : S_{\pm}^+ \rightarrow \mathcal{M}$, we first compute $\alpha, \beta \in \mathbb{N} \setminus \{0\}$ such that for all $m_1, m_2, m_3 \in \mathcal{M}$ and all $\tau \in \mathbb{N}$ $m_1 m_2^\beta m_3 = m_1 m_2^{\alpha\tau+\beta} m_3$ (it is an easy standard fact that such α, β exist and are computable). In particular, $u_1 u_2^\beta u_3 \in \mathcal{R}_e$ if and only if $u_1 u_2^{\alpha\tau+\beta} u_3 \in \mathcal{R}_e$. Then we modify the system Λ as follows: for each $A \in \mathcal{B}$ and $k \in \{1, \dots, n_A\}$ such that $t(Q_{A,k}) \leq \beta$, replace the variable $\tau_{A,k}$ by the constant $t(Q_{A,k})$; for each $t(Q_{A,k}) > \beta$, write by Euclidean division $t(Q_{A,k}) = \beta + \alpha t'_{A,k} + \varepsilon_{A,k}$ for some $\varepsilon_{A,k} < \alpha$ and $t'_{A,k} \in \mathbb{N}$; then replace the unknown $\tau_{A,k}$ by $\tau'_{A,k}$ using the identity $\tau_{A,k} = \alpha \tau'_{A,k} + \beta + \varepsilon_{A,k}$. We get a linear system of integer equations with unknowns $\tau'_{A,k}$, with at most $2\#\pi_0(D)$ equations, with coefficients bounded in terms of α and β , and with $l \leq 6\#\pi_0(D)$ unknowns. As above, one can compute a bound M' on all minimal solutions of all such systems which have a solution. The same argument as above shows that $\alpha M' + \beta$ is an upper bound the exponent of periodicity of the shortest solution of Σ , including rational constraints. $\square$

Remark. Using estimates on the size of minimal solutions of linear systems of Diophantine equations, and being more careful in the analysis, one can get an explicit bound of $M = 2^{O(\#\pi_0(D)) + n \log n}$ where n is the size of an automaton accepting the languages of $\underline{\mathcal{R}}$, see [DGH05].

5 Prelaminations

5.1 Definition

Definition 5.1. A prelamination $\mathcal{L}$ on a band complex Σ consists in

- A finite set $V(\mathcal{L}) \subset D$ containing all vertices of Σ . Points of $V(\mathcal{L})$ are called $\mathcal{L}$ -subdivision points.
- For each band $B = [a, b] \times [0, 1]$, a set $L_B(\mathcal{L}) \subset [a, b] \times [0, 1]$ of the form $\{c_1, \dots, c_r\} \times [0, 1]$. Each segment $\{c_i\} \times [0, 1]$ is called a leaf segment. The endpoints of each leaf segment are asked to lie in $V(\mathcal{L})$: $f_{B,\varepsilon}(c_i, \varepsilon) \in V(\mathcal{L})$ for $\varepsilon = 0, 1$. We also require that both segments $\{a\} \times [0, 1]$ and $\{b\} \times [0, 1]$ lie in $L_B(\mathcal{L})$.

See Figure 1 for an illustration.

Remark 5.2. In this definition, the precise value of the attaching maps $f_{B,\varepsilon}$ are not important, except at endpoints of leaf segments. Moreover, the precise position of the $\mathcal{L}$ -subdivision points in D is unimportant, only the induced ordering matters. Thus, a prelamination of a band complex is really determined by the ordering of the elements of $V(\mathcal{L})$ in each component of D , and for each band, by a bijection preserving or reversing the ordering, between a subset of $V(\mathcal{L}) \cap J_0$ and a subset of $V(\mathcal{L}) \cap J_1$.

We also view leaf segments of $\mathcal{L}$ as subsegments of the topological realisation $|\Sigma|$ of Σ . A *leaf* of $\mathcal{L}$ is a connected component of the union of the leaf segments in $|\Sigma|$. A *leaf path* is a concatenation of oriented leaf segments (which defines a path contained in a leaf).

A leaf segment $\{c\} \times [0, 1] \subset [a, b] \times [0, 1]$ is *singular* if $c \in \{a, b\}$. A leaf path is *regular* if it consists of non-singular leaf segments. A leaf is *singular* if it contains a singular leaf segment, i. e. if it contains a vertex of Σ .

An *elementary segment* of $\mathcal{L}$ is a segment of D joining two adjacent $\mathcal{L}$ -subdivision points. The set of oriented elementary segments of $\mathcal{L}$ is denoted by $E(\mathcal{L})$. We say that $J \subset D$ is *adapted to $\mathcal{L}$* if it is a union of elementary segments of $\mathcal{L}$.

A point $x \in V(\mathcal{L})$ is *not an orphan* if for every base J_ε of any band B , with $x \in J_\varepsilon$, there is a leaf segment of B having x as an endpoint: $x \in f_{B,\varepsilon}(L_B(\mathcal{L}))$. On Figure 1, orphans are represented as thick dots. The prelamination is *complete* if no $\mathcal{L}$ -subdivision point is an *orphan* (in this case, the prelamination could be called a genuine *lamination*).

If $B = [a, b] \times [0, 1]$ is a band with attaching maps $f_{B,0}, f_{B,1}$, we define the *opposite* B^{-1} of B as the band $[a, b] \times [0, 1]$ with attaching maps $f_{B^{-1},i} = f_{B,1-i}$ for $i = 0, 1$, with twisting morphism $\varphi_{B^{-1}} = \varphi_B^{-1}$. An *oriented band* is a band of Σ or its opposite. The leaf segments of B define leaf segments of B^{-1} . We orient each leaf segment $\{x\} \times [0, 1]$ of the oriented band B from $(x, 0)$ to $(x, 1)$ so that leaf segments of B and B^{-1} have the opposite orientation. If B is an oriented band with bases J_0, J_1 , we denote by $\text{dom } B = J_0$ its initial base, and we call it its *domain*.

A Σ -word is a word on the alphabet of oriented bands of Σ . The twisting morphism of the Σ -word $w = B_1 \dots B_n$ is $\varphi_w = \varphi_{B_1} \circ \dots \circ \varphi_{B_n}$. To each leaf path naturally corresponds a Σ -word corresponding to the oriented bands crossed by this path. We write $[x, w)$ for the leaf path starting at the point x and whose Σ -word is w (this leaf path is unique, but may fail to exist for some x and w). The *holonomy* of w is the map h_w defined on a subset of $V(\mathcal{L})$ and mapping x to the terminal endpoint of $[x, w)$ when $[x, w)$ exists. We write paths from right to left so that $h_{ww'} = h_w \circ h_{w'}$ (thus, the path ww' starts by w' and ends by w). By extension, if $[a, b] \subset D$ is such that h_w maps a to a' and b to b' , we say that h_w maps $[a, b]$ to $[a', b']$. When such w exists, we say that $[a, b]$ and $[a', b']$ are *equivalent under the holonomy of $\mathcal{L}$* .

5.2 Möbius strips

A *Möbius strip* S in a prelaminated band complex $(\Sigma, \mathcal{L})$ consists of a reduced band word w and an interval I adapted to $\mathcal{L}$ such that h_w maps I to itself reversing the orientation, and $h_{w_1}(I) \neq h_{w_2}(I)$ for any two distinct non-empty suffixes w_1, w_2 of w . Note that $(\Sigma, \mathcal{L})$ has only finitely many Möbius strips. A *core leaf* of S is a leaf path $[x, w)$ with $h_w(x) = x$.

Remark 5.3. Note that if the twisting morphism φ_w of w is trivial, or more specifically, if we are solving untwisted equations in a free group, the presence of a Möbius strip prevents the existence of a reduced solution of Σ . Thus, in the context of equations in free groups, prelaminations containing Möbius strips can be discarded.

If $\mathcal{L}$ is complete and all its Möbius strips have a core leaf, we say that $\mathcal{L}$ is *Möbius-complete*. A leaf is *pseudo-singular* if it is singular or if it contains the core leaf of a Möbius strip. Most prelaminations we will use will consist of pseudo-singular leaves.

If S has no core leaf, *adding a core leaf to S* means extending $\mathcal{L}$ by adding at most $|w|$ new leaf segments $l_1, \dots, l_n$ so that S has a core leaf l in the obtained prelamination, with $l \supset l_1 \cup \dots \cup l_n$.

5.3 Rational constraints on the prelamination

Consider a band complex Σ with rational constraints in standard form given by $\rho : S_{\pm}^+ \rightarrow \mathcal{M}$ and $\underline{m} \in \mathcal{M}^{E(\Sigma)}$. Let $\mathcal{L}$ be a prelamination on Σ . Recall that $E(\mathcal{L})$ is the set of oriented elementary segments of $\mathcal{L}$, i. e. of subsegments of D joining adjacent $\mathcal{L}$ -subdivision points.

A set of rational constraints on the prelamination $\mathcal{L}$ is a tuple $m'_e \in \mathcal{M}^{E(\mathcal{L})}$ such that $m'_e = \overline{m'_e}$ and which refines $\underline{m}$ in the following sense. For a segment J written as a concatenation of elementary segments of $\mathcal{L}$, $J = e_1 \dots e_n$, define $m'_J = m'_{e_1} \dots m'_{e_n} \in \mathcal{M}$. Then $\underline{m}'$ refines $\underline{m}$ if for all J oriented elementary segment of Σ , $m_J = m'_J$. This compatibility allows to blur the distinction between $\underline{m}$ and $\underline{m}'$, and we will use the same notation $\underline{m}$ for both.

5.4 Prelaminations and rational constraints induced by a solution

Consider a band complex Σ with $\rho : S_{\pm}^+ \rightarrow \mathcal{M}$ and $\underline{m} = (m_e)_{e \in E(\Sigma)}$ a set of rational constraints standard form. Let σ be a solution of $(\Sigma, \underline{m})$. The labelling σ allows to subdivide D into subsegments, each of which is labelled by a letter in $S_{\pm}$. This defines a complete prelamination $\mathcal{L}'$ whose subdivision points are the set of vertices of the subdivided D : since σ is a solution, each band B defines a pairing between all the vertices contained in its two bases. This pairing can be realised geometrically by a finite set of leaf segments of the form $\{c\} \times [0, 1] \subset [a, b] \times [0, 1]$ by changing the two attaching maps of B relatively to the endpoints of the bases.

Let $\mathcal{L}_{\infty}(\sigma) \subset \mathcal{L}'$ be the complete prelamination consisting of the pseudo-singular leaves of $\mathcal{L}'$. We call $\mathcal{L}_{\infty}(\sigma)$ the *complete prelamination induced by σ* . This is a Möbius-complete prelamination because the group Φ of twisting morphisms has no inversion.

For any segment J adapted to some prelamination $\mathcal{L} \subset \mathcal{L}_{\infty}(\sigma)$, one can define $m_J = \rho(\sigma_J) \in \mathcal{M}$, which defines a rational constraint on $\mathcal{L}$. This leads to the following definition:

Definition 5.4. A prelamination $\mathcal{L}$ with rational constraints $\mathcal{M}$ is induced by σ if $\mathcal{L} \subset \mathcal{L}_{\infty}(\sigma)$, all leaves of $\mathcal{L}$ are pseudo-singular, and if $m_e = \rho(\sigma_e)$ for all $e \in E(\mathcal{L})$.

Remark 5.5. The rational constraints induced by a solution are *invariant under the holonomy* in the following sense: consider a Σ -word w with twisting morphism φ_w whose holonomy maps some $e \in E(\mathcal{L})$ to $e' \in E(\mathcal{L})$, then $m_{e'} = \varphi_w(m_e)$.

Given a prelamination $\mathcal{L}$ of Σ , we define the solutions of $(\Sigma, \mathcal{L})$. A *labelling* σ of $(\Sigma, \mathcal{L})$ is a labelling of each $e \in E(\mathcal{L})$ by a word $\sigma_e \in S_{\pm}^+$ so that $\sigma_{\bar{e}} = \overline{\sigma_e}$. As in section 3.2, this allows to define $\sigma_J \in S_{\pm}^+$ for all oriented segment J adapted to $\mathcal{L}$.

Definition 5.6. A *solution* of $(\Sigma, \mathcal{L})$ is a labelling σ of $(\Sigma, \mathcal{L})$ such that for all Σ -word w and all oriented segments J, J' adapted to $\mathcal{L}$ such that the holonomy of w maps J to J' , then $\sigma_{J'} = \varphi_w(\sigma_J)$.

If there are rational constraints on $\mathcal{L}$, then we additionally require that $\rho(\sigma_e) = m_e$ for all $e \in E(\mathcal{L})$.

The following observation is obvious.

Lemma 5.7. *Let σ be a solution of Σ with rational constraints $\underline{m}$. Let $(\mathcal{L}, \underline{m}')$ be a prelamination with rational constraints induced by σ .*

Then σ is a solution of $(\Sigma, \mathcal{L})$ with rational constraints $\underline{m}'$. $\square$

Definition 5.8. *Consider $\mathcal{L} \subset \mathcal{L}'$ some prelaminations on Σ with rational constraints $\underline{m}, \underline{m}'$. We say that $\mathcal{L}'$ extends $\mathcal{L}$ and we denote $\mathcal{L} \prec \mathcal{L}'$ if for all $e \in E(\mathcal{L})$, $m_e = m'_e$.*

When $\mathcal{L}'$ extends $\mathcal{L}$, the set of solutions of $(\Sigma, \mathcal{L}')$ with constraints $\underline{m}'$ is a subset of the set of solutions of $(\Sigma, \mathcal{L})$ with constraints $\underline{m}$.

6 Algorithms on prelaminations

We now present two algorithms that will be part of the main algorithm solving equations. One of them enumerates prelaminations of a given a band complex. The other is given a prelamination, and checks several properties, for instance to prove that it cannot be induced by a shortest solution.

6.1 A first algorithm: the prelamination generator

We first consider a *prelamination generator*: this straightforward algorithm takes as input a band complex with rational constraints Σ and enumerates a set of prelaminations, such that any prelamination with rational constraints induced by a solution of Σ has an extension which is enumerated.

The set of produced prelaminations is organised in a locally finite rooted tree. The prelamination at depth 0 is the *root prelamination* whose leaf segments are the boundary segments of the bands. By definition, the root prelamination is contained in any prelamination. We define the rational constraints on the root prelamination as the rational constraints on Σ . The children of a given prelamination $\mathcal{L}$ are the extensions of $\mathcal{L}$ produced by the following extension process, consisting of the three following steps:

Step 1: extend leaves of the prelamination. Let $\mathcal{O}$ be the set of orphans of $\mathcal{L}$ (as defined in Section 5.1). Look at all the possible ways to add leaf segments to $\mathcal{L}$, so that every leaf segment added has an endpoint in $\mathcal{O}$, and points in $\mathcal{O}$ are no more orphans in the new prelamination.

Step 2: add cores of Möbius strips. Consider all the (finitely many) Möbius strips of $(\Sigma, \mathcal{L})$ having no core leaf. Then look at all possible ways to add a core leaf to each of them (see section 5.2 for definitions).

Step 3: extend rational constraints. If $\mathcal{L}'$ is a prelamination constructed above, and if $\mathcal{L}' = \mathcal{L}$, discard $\mathcal{L}'$. Given $\mathcal{L}' \supsetneq \mathcal{L}$ constructed above, and a set of rational constraints $\underline{m} \in \mathcal{M}^{E(\mathcal{L})}$ on $\mathcal{L}$, consider all possible rational constraints $\underline{m}' \in \mathcal{M}^{E(\mathcal{L}')}$ such that $(\mathcal{L}', \underline{m}')$ extends $(\mathcal{L}, \underline{m})$ as in Definition 5.8, meaning that for all oriented elementary segment $e \in E(\mathcal{L})$ written as a concatenation $e = e'_1 \dots e'_n$ of elements of $E(\mathcal{L}')$, $m_e = m'_{e'_1} \dots m'_{e'_n}$. Note that it might happen that for some $\mathcal{L}'$, there exists no set of rational constraints $\underline{m}'$.

Remark. The second step would actually be unnecessary if there was no twisting as no reduced solution of Σ can induce a prelamination containing a Möbius strip whose twisting morphism is trivial.

Step 2 should add a core leaf to every Möbius strip of $\mathcal{L}$, but it is allowed to create new Möbius strips having no core leaf.

If all rational constraints are given by one-letter constants, then the third step consists in ensuring that one does not subdivide the elementary segments corresponding to constants.

If $\mathcal{L}$ is Möbius-complete (i. e. has no orphan and all its Möbius strips have a core leaf), the only extension of $\mathcal{L}$ produced by the extension process is $\mathcal{L}$, and is discarded. In particular, $\mathcal{L}$ is a leaf of the rooted tree of prelaminations. If $\mathcal{L}$ is not Möbius-complete, all the prelaminations produced properly contain $\mathcal{L}$. Still, it may happen that no extension of $\mathcal{L}$ is produced if for all $\mathcal{L}' \supsetneq \mathcal{L}$ obtained after step 2, no extension of the rational constraints to $\mathcal{L}'$ exist. Of course, in this case, $\mathcal{L}$ cannot be induced by a solution of Σ .

Recall that all band complexes and prelaminations come with a set of rational constraints which we do not mention explicitly.

Lemma 6.1. *Consider σ a solution of Σ . Assume that σ induces some prelamination $\mathcal{L}$ produced by the prelamination generator.*

Then either $\mathcal{L} = \mathcal{L}_\infty(\sigma)$, or there exists some extension $\mathcal{L}'$ of $\mathcal{L}$ produced by the prelamination generator which is induced by σ .

Proof. Assume that $\mathcal{L} \subsetneq \mathcal{L}_\infty(\sigma)$. If $\mathcal{L}$ is not complete, then step 1 will clearly produce an extension of $\mathcal{L}_1 \supsetneq \mathcal{L}$ contained in $\mathcal{L}_\infty(\sigma)$. Otherwise, step 1 will produce only $\mathcal{L}_1 = \mathcal{L}$. Since all Möbius strips of $\mathcal{L}_\infty(\sigma)$ contain a core leaf, if $\mathcal{L}_1$ contains a Möbius strip without core leaf, then step 2 produces some $\mathcal{L}_2 \supsetneq \mathcal{L}_1$ contained in $\mathcal{L}_\infty(\sigma)$. The rational constraints induced by σ on $\mathcal{L}_2$ will be produced in step 3. The lemma follows. $\square$

Corollary 6.2. *For any solution σ of Σ , $\mathcal{L}_\infty(\sigma)$ will be produced by the prelamination generator.*

Consider some prelamination $\mathcal{L}$ which is not Möbius-complete, and such that none of its extensions constructed by the prelamination generator can be induced by a shortest solution of Σ . Then $\mathcal{L}$ cannot be induced by a shortest solution of Σ .

Proof. The first statement clearly follows from the lemma since $\mathcal{L}_\infty(\sigma)$ cannot contain an infinite chain of prelaminations strictly contained in each other. The second statement is clear. $\square$

6.2 A second algorithm: the prelamination analyser

We are going to describe a machine which analyses a given prelamination. We call it the *analyser*.

This algorithm takes as input a band complex Σ together with a prelamination $\mathcal{L}$ (with rational constraints). It tries to reject prelaminations for which it can prove that it cannot be induced by a shortest solution.

More precisely, if the input prelamination $\mathcal{L}$ is Möbius-complete, the analyser determines easily whether it is induced by some solution or not. The analyser then stops and outputs “Solution found” together with a solution, or “Reject” accordingly.

If the input prelamination $\mathcal{L}$ is not Möbius-complete, it looks for some certificate ensuring that no shortest solution can induce $\mathcal{L}$. If it can find one, it says “Reject” and stops. Otherwise, it may fail to stop or say “I don’t know”.

Actually, we could make sure that it actually always terminates because given $\mathcal{L}$, there are only finitely certificates to try. But we won’t need this fact.

The overall structure of the prelamination analyser is as follows.

Algorithm 6.3. Prelamination Analyser

Input: a prelamination with rational constraints $\mathcal{L}$ on a band complex Σ .

Output: “Solution found”, “Reject” or “I don’t know”.

- If $\mathcal{L}$ is a Möbius-complete prelamination, the algorithm stops and its output is either
 - “*Solution Found*”: the analyser can produce a solution inducing $\mathcal{L}$.
 - “*Reject*”: there exists no solution inducing $\mathcal{L}$, and the analyser can produce a certificate proving it.
- If $\mathcal{L}$ is not Möbius-complete, then the algorithm looks for certificates proving that $\mathcal{L}$ cannot be induced by a shortest solution of Σ . The algorithm tries four kinds of reasons to reject $\mathcal{L}$:
 - rejection for incompatibility of rational constraints (see subsection 6.2.1),
 - rejection for non-existence of invariant measure (see subsection 6.2.2),
 - rejection for too large exponent of periodicity (see subsection 6.2.3),
 - and rejection by a shortening sequence of moves (see subsection 6.2.4).

This procedure may stop or not, and its output can be

- “*Reject*”: the algorithm produces a certificate proving that $\mathcal{L}$ cannot be induced by a shortest solution of Σ .
- “*I don’t know*”: failure to find such a certificate.

We now describe more precisely the work of the prelamination analyser, describing the four types of rejection.

6.2.1 Incompatibility of rational constraints

The set of rational constraints $\underline{m}$ on a prelamination induced by a solution of Σ is invariant under the holonomy (see Remark 5.5): for all band B whose holonomy maps some $e \in E(\mathcal{L})$ to $e' \in E(\mathcal{L})$, $m_{e'} = \varphi_B(m_e)$. Thus, one can obviously reject rational constraints where this does not hold. For example, in this way one rejects prelaminations such that the constants do not match under the holonomy.

There is another source of incompatibility coming from the twisting morphisms. Assume that w is a Σ -word whose holonomy fixes some segment J adapted to $\mathcal{L}$, and whose twisting morphism φ_w is non-trivial. Then for any solution σ of Σ inducing $\mathcal{L}$, σ_J lies in $\text{Fix}\varphi_w$. By definition of standard form of rational constraints (section 3.5), $\text{Fix}\varphi_w$ is represented by $\rho : S_{\pm}^* \rightarrow \mathcal{M}$, so $\sigma_J \in \text{Fix}\varphi_w$ if and only if $\rho(\sigma_J) \in F$ for the subset $\rho(\text{Fix}\varphi_w) \subset \mathcal{M}$ which has already been computed. Thus, one can reject $\mathcal{L}$ if $m_J \notin \rho(\text{Fix}\varphi_w)$ for some w whose holonomy fixes J . Even though the set of Σ -words

whose holonomy fixes J may be infinite, this fact can be algorithmically checked. Indeed, the fact that $m_J \in \rho(\text{Fix}\varphi_w)$ needs to be checked only for a generating set of the group of Σ -words w fixing J , and such a generating set is easily computed.

We say that $\mathcal{L}$ shows an *incompatibility of rational constraints* when one can reject $\mathcal{L}$ for one of the two possible reasons above. In the prelamination analyser, rejection for incompatibility of rational constraints simply consists in checking this algorithmically.

When $\mathcal{L}$ is Möbius-complete, this is the only obstruction to the existence a solution of Σ inducing $\mathcal{L}$ and $\underline{m}$:

Lemma 6.4. *Let $\mathcal{L}$ be a Möbius-complete prelamination of Σ . Then Σ has a solution inducing $\mathcal{L}$ with set of rational constraints $\underline{m}$ if and only if it shows no incompatibility of rational constraints.*

This explains how the prelamination analyser determines the existence of a solution when its input is a Möbius-complete prelamination: it just checks whether $\mathcal{L}$ shows an incompatibility of rational constraints. The argument below also shows how to produce a solution in the absence incompatibility of rational constraints.

Proof. Since $\mathcal{L}$ is Möbius-complete, the set of oriented elementary segments $E(\mathcal{L})$ is partitioned into orbits under the holonomy of $\mathcal{L}$, and no $e \in E(\mathcal{L})$ is the orbit of $\bar{e}$ (i. e. e with the orientation reversed). For each such orbit, choose one representative $e \in E(\mathcal{L})$, and choose some word $\sigma_e \in \rho^{-1}(m_e)$ (ρ is onto). For each e' in the orbit of e , choose a Σ -word w such that $h_w(e) = e'$ and define $\sigma_{e'} = \varphi_w(\sigma_e)$. Since $\mathcal{L}$ shows no incompatibility of rational constraints of the first kind, one has $\rho(\sigma_{e'}) = \varphi_w(m_e) = m'_e$. If w' is another word with $h_{w'}(e) = e'$, then $\sigma_e \in \text{Fix}\varphi_{w'w^{-1}}$ because $\mathcal{L}$ shows no incompatibility of rational constraints of the second kind, so $\sigma_{e'}$ does not depend on the choice of w' . Since e and $\bar{e}$ are not in the same orbit, we can extend these choices by defining $\sigma_{\bar{e}'} = \overline{\sigma_{e'}}$ for all e' in the orbit of e . Doing this for every orbit of elementary segments, we get a solution of $(\Sigma, \mathcal{L})$ respecting the rational constraints. $\square$

6.2.2 Rejection for non-existence of invariant measure

If a prelamination $\mathcal{L}$ is induced by a solution σ , then the labelling defines a positive length l_e on each elementary segment e of $\mathcal{L}$, and more generally, to each segment adapted to $\mathcal{L}$. This can be viewed as a combinatorial measure invariant under the holonomy along the leaves of $\mathcal{L}$ in the following sense: if I and J are segments adapted to $\mathcal{L}$ and such that the holonomy h_B maps I to J for some band B , then $l_I = l_J$.

A prelamination may fail to have such an invariant transverse measure, for example if the holonomy along the leaves maps a segment to a proper subset of itself. The existence a combinatorial measure (l_e) is equivalent to the existence of a positive solution to some system of linear equations with integer coefficients, which can be checked algorithmically. Thus, the prelamination analyser can check whether $\mathcal{L}$ admits an invariant combinatorial measure, and if there is no such measure, it rejects $\mathcal{L}$.

6.2.3 Rejection for too large exponent of periodicity

In some cases, one can read in a prelamination $\mathcal{L}$ the fact that any solution inducing $\mathcal{L}$ must have a very large exponent of periodicity.

Definition 6.5. *We say that $\text{exponent}(\mathcal{L}) \geq N$ if there exists a segment J which is a concatenation of N intervals $I_1 \cdots I_N \subset D$ adapted to $\mathcal{L}$, and some Σ -words w_i whose holonomy maps I_i to I_{i+1} preserving the orientation, and whose twisting automorphisms φ_{w_i} are trivial.*

Clearly, if $\text{exponent}(\mathcal{L}) \geq N$, then the exponent of periodicity of any solution inducing $\mathcal{L}$ is at least N . By Bulitko's Lemma, one can compute a bound B such that any shortest solution of Σ has an exponent of periodicity bounded by B .

In the prelamination analyser, rejection for too large exponent of periodicity consists in computing B , and checking whether $\text{exponent}(\mathcal{L}) > B$. This is easily done algorithmically.

6.2.4 Rejection by a shortening sequence of moves

This last type of rejection is more complicated. In Section 7.2, we will introduce a finite set of moves which can be performed on a band complex with a prelamination. There will be several types of moves.

Definition 6.6 (Inert move). *We say that a prelaminated band complex $(\Sigma', \mathcal{L}')$ is obtained by an inert move from $(\Sigma, \mathcal{L})$ if*

- *for any solution σ of Σ inducing $\mathcal{L}$, there is a solution σ' of Σ' inducing $\mathcal{L}'$ such that $|\sigma'| = |\sigma|$*
- *for any solution σ' of Σ' (which need not induce $\mathcal{L}'$), there is a solution σ of Σ (which need not induce $\mathcal{L}$) and such that $|\sigma| = |\sigma'|$*

An obvious property of an inert move is that if $\mathcal{L}$ is induced by a shortest solution of Σ , then $\mathcal{L}'$ is induced by a shortest solution of Σ' . Examples of inert moves include band subdivision, domain cut, and band removal moves (see section 7.2).

To illustrate the two other types of moves, we start with an example: *the pruning move*. The input is a band complex Σ with domain D together with a prelamination $\mathcal{L}$, and the output is a new prelaminated band complex $(\Sigma', \mathcal{L}')$. We assume that there is a segment $[a, b] \subset D$ where a, b are vertices of Σ , such that $[a, b]$ is contained in a base J_0 of some band B , and such that no other base of Σ intersects the interior of $[a, b]$ (see Figure 4). Assume that a and b are not orphans in $\mathcal{L}$. The rational constraints on $(\Sigma, \mathcal{L})$ are assumed to be invariant under the holonomy. We view the band B as $J_0 \times [0, 1]$, the attaching map $f_{B,0}$ being the identity. Let $D' = D \setminus (a, b)$, and let Σ' be the band complex with domain D' obtained by replacing the band $B = J_0 \times [0, 1]$ by the two bands defined by the connected components of $(J_0 \setminus (a, b)) \times [0, 1]$. If a or b is an endpoint of D , then (a, b) has to be interpreted as the interior of $[a, b]$ in D , in which case $(J_0 \setminus (a, b)) \times [0, 1]$ consists of only one band or no band at all. The attaching map of the new bands are well defined because a and b are not orphans. The new prelamination is the one naturally obtained by restriction. The new set of rational constraints on $\mathcal{L}'$ is obtained by restriction. The set rational constraints $\underline{m}'$ on Σ' is the one induced by $\mathcal{L}'$: any elementary segment




Figure 4: Pruning move

e' of Σ' is a concatenation of elementary segments $e'_1, \dots, e'_n$ of $\mathcal{L}'$, which allows to define $m'_{e'} = m'_{e'_1} \dots m'_{e'_n}$.

It is clear that any solution σ of Σ inducing $\mathcal{L}$ (together with its set of rational constraints, which we don't mention explicitly) defines a solution σ' of Σ' inducing $\mathcal{L}'$ (with rational constraints). Moreover, the length of σ' is the length of the restriction of σ to D' .

Conversely, if σ' is any solution of Σ' , without assuming that σ' induces $\mathcal{L}'$, it naturally extends to a solution σ of Σ , and $|\sigma| \leq 2|\sigma'|$. We say that the move from $(\Sigma, \mathcal{L})$ to $(\Sigma', \mathcal{L}')$ is a *restriction move* and that the move from Σ' to Σ is an *extension move*.

Remark 6.7. Note that restriction moves are about prelaminated band complexes while extension moves are about band complexes without prelamination.

We will consider long sequences of restriction and extension moves which may leave a large part of the band complex essentially untouched. To keep track of this, we partition the domain D of the involved band complexes into $D = D_I \sqcup D_A$ where D_I and D_A are a union of connected components of D where D_I is the *inert* part (which is left untouched by the move), and D_A the *active* part. If D' is the domain of Σ' , we denote by $D'_I \sqcup D'_A$ the corresponding decomposition of D .

Definition 6.8 (Restriction move). *We say that $(\Sigma', \mathcal{L}')$ is obtained by a restriction move from $(\Sigma, \mathcal{L})$ if there is an injective map $\iota : D' \hookrightarrow D$ with $D_I = \iota(D'_I)$ and $\iota(D'_A) \subset D_A$ adapted to $\mathcal{L}$, such that for any solution σ of Σ inducing $\mathcal{L}$, there is a solution σ' of Σ' inducing $\mathcal{L}'$ such that*

- $|\sigma'_{D'_I}| = |\sigma_{D_I}|$
- $|\sigma'_{D'_A}| = |\sigma_{\iota(D'_A)}|$

Remark 6.9. We call ι the *restriction map*. For the pruning move, for instance, ι is just the inclusion.

Definition 6.10 (Extension move). *We say that Σ is obtained from Σ' by an extension move with Lipschitz factor $\lambda \geq 1$ if for any solution σ' of Σ' , there is a solution σ of Σ such that*

- $|\sigma_{D_I}| = |\sigma'_{D'_I}|$
- $|\sigma_{D_A}| \leq \lambda |\sigma'_{D'_A}|$.

For instance, one can take $\lambda = 2$ for the pruning move, under the requirement that D_A contain both bases of the band pruned.

To be able to ensure that a prelamination $\mathcal{L}$ cannot be induced by a shortest solution, we need a way of certifying that some subset $\iota(D'_A) \subset D_A$ is *short relative to D_A* , i. e. that for any solution σ inducing $\mathcal{L}$, $|\sigma|_{D'_A}|$ is at most $\varepsilon|\sigma|_{D_A}|$ for some small $\varepsilon > 0$. Relative shortness will be guaranteed by some repetition as in the lemma below.

Definition 6.11 (Certificate of shortness of J relative to D_A). *Let $(\Sigma, \mathcal{L})$ be a prelaminated band complex, and $J \subset D_A$ be a subset adapted to $\mathcal{L}$. A certificate of ε -shortness of J relative to D_A is a family of adapted subsets $J_1, \dots, J_N \subset D_A$ with disjoint interiors for some $N \geq 1/\varepsilon$, and which are all equivalent to J under the holonomy of $\mathcal{L}$.*

Lemma 6.12. *If there is a certificate of ε -shortness of J relative to D_A , then for any solution σ of Σ inducing $\mathcal{L}$, then $|\sigma|_J| \leq \varepsilon|\sigma|_{D_A}|$.*

Proof. This is essentially obvious: for any solution σ inducing $\mathcal{L}$, the labellings of $J_1, \dots, J_N$ will have the same length as J . Since the J_i 's don't overlap and are contained in D_A , we get $N \cdot |\sigma|_J| \leq |\sigma|_{D_A}|$ and the lemma follows. $\square$

The prelamination analyser will try to apply a sequence of elementary moves (as defined in Section 7.2) to the prelaminated band complex. It is clear that a succession of inert moves defines an inert move. Moreover, a succession of restriction moves having the same inert part also defines a restriction move, whose restriction map ι is the composition of the restriction maps of the elementary moves. Similarly, a succession of elementary extension moves having the same inert part defines an extension move whose Lipschitz factor is the product of the Lipschitz factors of the individual moves.

Definition 6.13 (Shortening sequence of moves). *A shortening sequence of moves for the prelamination $\mathcal{L}_0$ of Σ_0 consists of the following data:*

1. *a sequence of inert moves transforming $(\Sigma_0, \mathcal{L}_0)$ to $(\Sigma, \mathcal{L})$;*
2. *a sequence of elementary restriction moves, whose concatenation defines a restriction move $(\Sigma, \mathcal{L}) \rightarrow (\Sigma', \mathcal{L}')$, with corresponding map $\iota : D'_A \rightarrow D_A$;*
3. *a sequence of elementary extension moves transforming Σ' back to Σ ;*
4. *a certificate of ε -shortness of $\iota(D'_A)$ relative to D_A with $\varepsilon < 1/\lambda$, where λ is the product of the Lipschitz factors of the extension moves used above.*

Lemma 6.14. *If there exists a shortening sequence of moves for $\mathcal{L}_0$, then $\mathcal{L}_0$ cannot be induced by a shortest solution of Σ_0 .*

Proof. Assume on the contrary that $\mathcal{L}_0$ is induced by a shortest solution of Σ_0 . Using the property of inert moves, $\mathcal{L}$ is itself induced by a shortest solution σ of Σ . Using the restriction move, consider σ' a solution of Σ' such that $|\sigma'_{D'_I}| = |\sigma_{D_I}|$ and $|\sigma'_{D'_A}| = |\sigma_{\iota(D'_A)}|$. Using the extension move, consider a solution $\tilde{\sigma}$ of Σ such that $|\tilde{\sigma}_{D_I}| = |\sigma'_{D'_I}|$ and $|\tilde{\sigma}_{D_A}| \leq \lambda|\sigma'_{D'_A}|$. Since $\iota(D'_A)$ has a certificate of ε -shortness relative to D_A , $|\tilde{\sigma}| \leq |\sigma_{D_I}| + \varepsilon\lambda|\sigma_{D_A}| < |\sigma|$, a contradiction. $\square$

6.2.5 More about the analyser

We are now ready to describe the rejection by shortening moves by the prelamination analyser. The analyser tries all possible sequences of moves looking like a shortening sequence of moves constructed as follows: a sequence of inert moves transforming $(\Sigma_0, \mathcal{L}_0)$ into some $(\Sigma, \mathcal{L})$, followed by a choice of decomposition $D = D_I \sqcup D_A$, followed by a sequences of shortening moves with inert part D_I transforming $(\Sigma, \mathcal{L})$ into some $(\Sigma', \mathcal{L}')$, followed by a sequence of extension moves with inert part D_I and transforming Σ' back into Σ . All such sequences can be enumerated. Then look for $1/\lambda$ -shortness certificates for $\iota(D'_A)$ in D_A . Being a little more careful, one could easily ensure that this enumeration terminates, but we won't need this fact. If such a certificate exists, it will be found. In this case, the prelamination analyser rejects $\mathcal{L}$.

Lemma 6.15. *The prelamination analyser is correct: if it says “Solution found”, then Σ has a solution inducing $\mathcal{L}$; if it says “Reject”, then $\mathcal{L}$ cannot be induced by a shortest solution.*

Proof. The correctness when a solution has been found follows from Lemma 6.4. The correctness for each reason of rejection follows from Subsections 6.2.1, 6.2.2, 6.2.3, and from Lemma 6.14 above. $\square$

7 Asymptotic of prelaminations: Rips band complexes

In this section, we assume that in the locally finite tree of prelaminations produced by the prelaminations generator, there is an infinite chain, i. e. an infinite sequence of prelaminations with rational constraints extending each other.

By passing to a limit, we will construct a topological foliation on Σ together with an invariant transverse measure μ . With an assumption on the exponent of periodicity, we will prove that μ has no atom. However, μ may fail to have full support. Collapsing segments of measure zero, we will get a new foliated band complex Σ_μ where the measure is the Lebesgue measure: we say that Σ_μ is a *Rips band complex*.

In [GLP94, BF95] is studied the decomposition of such an object into so-called minimal components, and the classification of such minimal components. In particular, there are so-called homogeneous components, in which, as we will see, one can find arbitrarily high exponent of periodicity. If a minimal component is not homogeneous, one can perform a sequence of moves on the Rips band complex (to obtain so-called independent bands) so that it becomes either a surface component (also called interval exchange component) or an exotic component (also called thin, or Levitt). In both cases, we will prove the existence of a shortening sequence of moves (in the sense of Definition 6.13) for all sufficiently large prelaminations in our infinite chain.

These properties are used to prove that the main algorithm 8.2 always terminates, see Section 8.

7.1 Limit measured foliation

Consider an infinite sequence of prelaminations with rational constraints extending each other: $\mathcal{L}_1 \prec \mathcal{L}_2 \prec \dots$ (Definition 5.8).

The goal of this subsection is to associate to such a chain of prelaminations a topological foliation on Σ , and a transverse invariant measure under the assumption that each $\mathcal{L}_i$ has an invariant transverse measure (Subsection 6.2.2). Of course, those objects are not accessible to the algorithm, but they are used to prove the existence of certificates rejecting $\mathcal{L}_i$ for i large enough.

7.1.1 The limit topological foliation

We first define the notion of a topological foliation on a band complex Σ . Recall that bands of a band complex are defined by attaching maps $f_{B,\varepsilon} : [a, b] \times \{\varepsilon\} \rightarrow D$ for $\varepsilon = 0, 1$, but the only relevant data is the combinatorial arrangement of the endpoints of bases of bands. If Σ is endowed with a prelamination, more data has to be extracted from f_B , i. e. the combinatorial arrangement of the images of endpoints of leaf segments. A *topological foliation* $\mathcal{F}$ consists in choosing for each band, an injective continuous map $\tilde{f}_{B,\varepsilon} : [a, b] \times \{\varepsilon\} \rightarrow D$ coinciding with $f_{B,\varepsilon}$ on endpoints of bases of bands. Here, the precise values of $\tilde{f}_{B,\varepsilon}$ matters, not only its values on a finite subset.

Leaf segments of $\mathcal{F}$ in a band B consist of all segments $\{x\} \times [0, 1] \subset B$. The *endpoints* of the leaf segments are the points $f_{B,0}(x, 0), f_{B,1}(x, 1) \in D$. Consider the equivalence relation on the set of leaf segments, generated by the property of having a common endpoint. A *leaf* of $\mathcal{F}$ is the union of leaf segments in a given equivalence class. As usual, we identify a leaf with its image in the topological realisation of the band complex.

As for prelaminations, we can talk about singular or regular leaf segments, leaf paths, and leaves. Given a Σ -word w , and $x \in D$, we still use the notation $[x, w)$ which has a meaning for a topological foliation. The holonomy map h_w is now a homeomorphism between a compact interval $\text{dom } w \subset D$, called the *domain* of w , and its image.

Definition 7.1. Consider a (maybe infinite) set $l_1, \dots, l_p, \dots$ of leaves of a topological foliation $\mathcal{F}$ containing all singular leaves.

We say that a sequence of prelaminations $\mathcal{L}_i$ is represented in $\mathcal{F}$ if one can write $\mathcal{L}_i$ as a union of leaf segments of $\mathcal{F}$ so that

- $\mathcal{L}_1 \subset \mathcal{L}_2 \subset \dots$ is an exhaustion of $l_1 \cup \dots \cup l_p \cup \dots$ by finite subgraphs,
- every Möbius strip of $\mathcal{L}_i$ has a core in $\mathcal{L}_j$ for some $j \geq i$.

The leaves l_i are called the *special leaves* of $\mathcal{F}$.

Remark 7.2. By definition, every singular leaf l is special. If l is pseudo-singular but not singular, and if $\mathcal{L}_i$ has an invariant combinatorial measure, then l is special. Indeed, consider $x \in \mathcal{L}$ and a Σ -word w such that x lies in the interior of $\text{dom } w$, $h_w(x) = x$, and w reverses the orientation. Let $[a, b]$ be the domain of h_{w^2} . If h_{w^2} fixes a , then there will a Möbius strip in $\mathcal{L}_i$ for i large enough since the leaf through a is singular hence special. The second condition ensures that l is special. Otherwise, up to changing w to w^{-1} , we can assume $h_{w^2}(a) > a$. This implies $h_{w^2}(b') < b'$, where $b' = h_w(a)$. It follows that the holonomy of w^2 maps $[a, b']$ to a proper subsegment of itself, which prevents the existence of an invariant combinatorial measure.

Proposition 7.3. *Consider an infinite sequence of prelaminations $\mathcal{L}_1 \prec \mathcal{L}_2 \prec \dots$ extending each other, and consisting of pseudo-singular leaves. Assume that for each subdivision point x of $\mathcal{L}_i$, there exists $j > i$ such that x is not orphan in $\mathcal{L}_j$, and that any Möbius strip of $\mathcal{L}_i$ has a core in some $\mathcal{L}_j$ for some $j \geq i$.*

Then there exists a topological foliation $\mathcal{F}$ on Σ representing $\mathcal{L}_i$.

Proof. Embed D in $\mathbb{R}$ and endow it with the induced metric. Denote by $|C|$ the diameter of a subset $C \subset D$. Let $L_i = V(\mathcal{L}_i)$ be the set of subdivision points of $\mathcal{L}_i$. Recall that we have some freedom in the choice of the points of L_i as long as we don't change the induced ordering of the points. Thus, we can choose inductively L_i so that $L_i \supset L_{i-1}$, and so that the following holds. For each component C of $D \setminus L_i$, let C' be the component of $D \setminus L_{i-1}$ containing C . Then either $C = C'$, or $|C| \leq |C'|/2$.

We also have some freedom in the choice of attaching maps of each band $f_{B,\varepsilon}^{(i)} : [a, b] \times \{0, 1\} \rightarrow D$ used to define the prelamination $\mathcal{L}_i$. By modifying $f_{B,\varepsilon}^{(i)}$ with respect to L_i , we can assume that $f_{B,\varepsilon}^{(i)}$ coincides with $f_{B,\varepsilon}^{(i-1)}$ on each C such that $C = C'$ (formally, we should rather say that they coincide on $f_{B,\varepsilon}^{-1}(C)$, but to keep notations lighter, we will make this abuse of notations).

Let $L_\infty = \bigcup_{i \geq 0} L_i$. Our inductive choice of L_i ensures that for any connected component C of $D \setminus \overline{L_\infty}$, its endpoints lie in some L_{i_0} so C is actually a connected component of $D \setminus L_{i_0}$. Say that a connected component of $D \setminus L_{i_0}$ is *stabilised* if it is also a connected component of $D \setminus \overline{L_\infty}$. In this case, any oriented band B whose initial base intersects (and therefore contains) C , the restriction to C of the attaching map $f_{B,0}^{(i)}$ is independent of $i \geq i_0$. Up to enlarging i_0 , we can assume that the endpoints of C are not orphan in $\mathcal{L}_{i_0}$. It follows that the image C' of C under the holonomy of B is also a stabilised component, so the restriction to C of the holonomy of B does not depend on $i \geq i_0$.

We fix a band B with bases $J_0, J_1 \subset D$. We claim that $h_i = f_{B,1}^{(i)}(f_{B,0}^{(i)})^{-1} : J_0 \rightarrow J_1$ converges uniformly as i goes to infinity. Fix some $\varepsilon > 0$. Let U be the (finite) union of all connected components of $D \setminus \overline{L_\infty}$ of diameter at least $\varepsilon/2$. Denote by $N_i \subset L_i$ the set of non-orphan subdivision points of $\mathcal{L}_i$, and note that $\bigcup_{i > 0} N_i = \bigcup_{i > 0} L_i$. Let i_0 be such that $\partial U \subset N_{i_0}$, and that all connected components of $(D \setminus U) \setminus N_{i_0}$ have diameter at most ε .

Consider $x \in J_0$. If $x \in U$, then $h_j(x) = h_i(x)$ for all $j, i > i_0$. If $x \notin U$, let $[s, s']$ be the smallest interval containing x such that s and s' are the endpoints of leaf segments of B . Then $h_i(s)$ and $h_i(s')$ are independent of i , and are at distance at most ε by choice of i_0 . Since for all i , $h_i(x) \in [h_i(s), h_i(s')]$, we get $|h_i(x) - h_j(x)| \leq \varepsilon$ for all $i, j \geq i_0$, which proves uniform convergence to some h_∞ . Since h_i^{-1} converges by the symmetric argument, h_∞ is a homeomorphism.

Finally, we can define the foliation using the attaching maps $\tilde{f}_{B,\varepsilon}$ defined by $\tilde{f}_{B,0} = f_{B,0}^{(0)}$ and $\tilde{f}_{B,1} = h_\infty \circ f_{B,0}^{(0)}$.

It is now clear that $\mathcal{F}$ represents $\mathcal{L}_i$: indeed $\mathcal{L}_i$ is an exhaustion of the union of the leaves it intersects because no subdivision point of $\mathcal{L}_i$ remains orphan forever. $\square$

Remark 7.4. Although the foliation is meaningless in the complement Σ^* of closure of special leaves, it can be convenient to ensure that the foliation on Σ^* is a twisted product, in other words, that the holonomy of any Σ -word preserving a connected component J of $\Sigma^* \cap D$ and preserving the orientation has to be the identity on J . This is easily done by choosing the attaching maps in the proof above to be affine on

the complement of the subdivision points. Also note that the measure to be defined in next section is zero on Σ^* .

7.1.2 The limit invariant measure

Proposition 7.5. *Consider an infinite sequence of prelaminations $\mathcal{L}_1 \prec \mathcal{L}_2 \prec \dots$ represented in some topological foliation $\mathcal{F}$. Assume that for each i , $\mathcal{L}_i$ has an invariant combinatorial measure.*

Then either the exponent of periodicity of $\mathcal{L}_i$ goes to infinity (see Definition 6.5), or there exists a measure μ on D , which has no atom and which is invariant under the holonomy of $\mathcal{F}$.

In particular, if no $\mathcal{L}_i$ is rejected by the prelamination analyser, then $\mathcal{F}$ has a non-atomic transverse invariant measure.

Proof. The construction follows [Pla75]. Denote by $\mathcal{F}^s$ the (finite) union of singular leaves of $\mathcal{F}$. We first note that if $\mathcal{F}^s$ is finite (i. e. contains only finitely many leaf segments), then $\mathcal{L}_i$ is eventually constant. Indeed, consider $J \subset D$ a segment meeting the singular leaves of $\mathcal{F}$ only at its endpoints. Since $\mathcal{L}_i$ preserves a combinatorial measure, the union of all non-singular pseudo-singular leaves of $\mathcal{L}_i$ intersects J in at most one point (its midpoint). Since all leaves of $\mathcal{L}_i$ are pseudo-singular, this bounds the number of leaf segments in $\mathcal{L}_i$.

Consider l_0 an infinite singular leaf of $\mathcal{F}$, and choose a base point $x_0 \in l_0$. Let $l_0(n)$ be the ball of radius n centred at x_0 in l_0 viewed as a graph, and consider $L_n = l_0(n) \cap D$.

We claim that $\#L_n$ grows polynomially with n :

Lemma 7.6. $\#L_n \leq 2(2n+1)^b$ where b is the number of bands of Σ .

Proof. Fix some n , and consider i large enough so that the finite graphs $l_0(n)$ is contained in the prelamination $\mathcal{L}_i$. Since $\mathcal{L}_i$ has an invariant combinatorial measure, one can embed D in $\mathbb{R}$ isometrically with respect to this combinatorial measure. For each band B of Σ , the holonomy of B extends to an isometry γ_B of $\mathbb{R}$. Let $G < \text{Isom}(\mathbb{R})$ be the subgroup generated by the isometries γ_B . One easily checks that the ball of radius n of the Cayley graph of G has cardinal $2(2n+1)^b$ where b is the number of bands of Σ . It follows that the set of images of x_0 under elements of length at most n of G has cardinal at most $2(2n+1)^b$. Since L_n is contained in this set, the lemma follows. $\square$

Consider the normalised counting measure $\mu_n = \frac{1}{\#L_n} \sum_{s \in L_n} \delta_s$ where δ_s is the Dirac mass at s . Consider the set $O_n = L_n \setminus L_{n-1}$. This is the locus where μ_n might fail to be preserved: if B is a band with bases J_0, J_1 , and if $x \in J_0 \cap L_n \setminus O_n$, then $h_B(x) \in L_n$ so $\mu_n(x) = \mu_n(h_B(x)) > 0$. Using the same argument for B^{-1} , this means that h_B preserves the measure μ_n except maybe at points of $O_n \cup h_B(O_n)$. Because of polynomial growth, there exists a subsequence n_k such that the proportion $\frac{\#O_{n_k}}{\#L_{n_k}}$ converges to 0 as k goes to ∞ .

Let μ be any accumulation point of μ_{n_k} for the weak-* topology (existence is ensured by Banach-Alaoglu's theorem). If we know that the endpoints of the bases of bands have zero measure, the following lemma proves that μ is invariant under the holonomy.

Lemma 7.7. *Consider a band B with bases J, J' and with holonomy $h : J \rightarrow J'$. Then $\mu|_{J'} = h_*\mu|_{\mathring{J}}$.*

Proof. Consider $\varepsilon > 0$ and a continuous function $f : D \rightarrow \mathbb{R}$ with support in $\mathring{J}$, and $f' = f \circ h^{-1}$ with support in $\mathring{J}'$. Then for all $\varepsilon > 0$, for k large enough,

$$\left| \int f' d\mu - \int f d\mu \right| \leq \frac{1}{\#L_{n_k}} \left| \sum_{s' \in L_{n_k}} f'(s') - \sum_{s \in L_{n_k}} f(s) \right| + \varepsilon.$$

For each $s \in L_{n_k} \setminus O_{n_k} \cap \mathring{J}$, the point $s' = h(s)$ lies in $L_{n_k} \cap \mathring{J}'$, and the two corresponding terms $f(s)$ and $f'(s')$ cancel. Thus,

$$\begin{aligned} \left| \int f' d\mu - \int f d\mu \right| &\leq \frac{1}{\#L_{n_k}} \sum_{s' \in O_{n_k}} |f'(s')| + \frac{1}{\#L_{n_k}} \sum_{s' \in O_{n_k}} |f(s)| + \varepsilon \\ &\leq \frac{\#O_{n_k}}{\#L_{n_k}} \|f\|_\infty + \varepsilon \end{aligned}$$

Since the proportion $\frac{\#O_{n_k}}{\#L_{n_k}} \rightarrow 0$ converges to zero, the lemma follows. $\square$

To prove that μ is invariant under the holonomy, we need to prove that μ has no atom. Note that the lemma implies that if $\mu(\{x\}) > 0$ for some $x \in \mathring{J}$, then $\mu(\{h(x)\}) = \mu(\{x\})$.

Consider $x \in D$ and a segment $J = [x, a]$. Consider the limit μ_J of the restriction of μ_{n_k} to J (up to passing to a subsequence). Clearly, μ_J coincides with μ on $\mathring{J}$, but $\mu_J(\{x\})$ may differ from $\mu(\{x\})$ because leaves of $\mathcal{L}_i$ can go in the neighbourhood of x without entering J . If $J' = [x, b'] \subset J$, then $\mu_{J'}(\{x\}) = \mu_J(\{x\})$. We thus denote by $\mu_+(x) = \mu_{[x, x+t]}(\{x\})$ and $\mu_-(x) = \mu_{[x-t, x]}(\{x\})$ for some (any) $t > 0$. Since $\mu_n(\{x\}) \rightarrow 0$ as $n \rightarrow \infty$, one gets $\mu_+(x) + \mu_-(x) = \mu(\{x\})$. The proof of Lemma 7.7 directly extends to the following:

Lemma 7.8. *Let J be a base of some band B with holonomy $h : J \rightarrow J'$, and assume that $[x, x + \varepsilon t] \subset J$ for some $\varepsilon = \pm 1$, $t > 0$. Let $\varepsilon' = \pm 1$ be the sign of $h(x + \varepsilon t) - h(x)$ for $t > 0$ small enough.*

Then $\mu_{\varepsilon'}(h(x)) = \mu_\varepsilon(x)$. $\square$

The following lemma ends the proof of Proposition 7.5. $\square$

Lemma 7.9. *If μ has an atom, then $\text{exponent}(\mathcal{L}_i) \rightarrow \infty$.*

Proof. Consider $a \in D$ with $\mu(\{a\}) > 0$. Assume for instance that $\mu_+(a) > 0$.

Let l be the $\mathcal{F}$ -leaf through a . Define the graph $\mathcal{C}$ as follows: its vertex set $V(\mathcal{C}) \subset (l \cap D) \times \{+1, -1\}$ is defined by $(y, \varepsilon) \in V(\mathcal{C})$ if there exists a Σ -word w whose holonomy h_w is defined on $[a, a + t]$ for some $t > 0$, with $h_w(a) = y$ and $h_w(a + t) - h_w(a)$ of the sign of ε . Put an edge labelled B between (y, ε) and (y', ε') if $h_B(y) = y'$ and $h_B(y + \varepsilon t) - h_B(y)$ is defined and of the sign of ε' for $t > 0$ small enough. The natural map $V(\mathcal{C}) \rightarrow l$ sending (y, ε) to y extends to a map $\mathcal{C} \rightarrow l$ sending edge to edge, and which is at most 2 to 1. Let us choose the point $(a, +1)$ (corresponding to the empty word) as a base point for $\mathcal{C}$.

By Lemma 7.8, for all $(y, \varepsilon) \in \mathcal{C}$, $\mu_\varepsilon(y) = \mu_+(a)$. Since $\mu(D) < \infty$, $\mathcal{C}$ is a finite graph. Since $\mu_+(a) > 0$, the singular leaf l_0 of $\mathcal{F}$ accumulates on a on the right.

Let T be a maximal tree in $\mathcal{C}$. For any vertex $(y, \epsilon) \in \mathcal{C}$, let us denote $[(a, +1), (y, \epsilon)]_T$ the unique path in T joining the two points. Each such segment defines a Σ -word $w_{(y, \epsilon)}$ whose domain contains an interval $[a, a + \eta_y]$ for some $\eta_y > 0$. For each oriented edge $e \in \mathcal{C}$ not in T , labelled by B , joining v_1 to v_2 , the Σ -word $w_e = w_{v_2}^{-1} B w_{v_1}$ has its domain containing some interval $[a, a + \eta_e]$ for some $\eta_e > 0$. Note that by definition of $\mathcal{C}$, its holonomy fixes a and preserves the orientation.

Let $\eta > 0$ be such that the holonomy of each Σ -word $w_e, e \in \mathcal{T} \setminus T$ is defined on $[a, a + \eta]$.

We claim that for any $x \in l_0 \cap D$ close enough to a on the right of a , there exists an oriented edge $e \in \mathcal{C} \setminus T$ such that $h_{w_e}([a, x]) \subsetneq [a, x]$. Otherwise, take x close enough to a so that the set $\bigcup_{v \in V(\mathcal{C})} h_{w_v}([a, x])$ does not contain an endpoint of a base of band. Then any band defined on some $h_{w_v}(x)$ is defined on the whole interval $h_{w_v}([a, x])$. If for all oriented edge $e \in \mathcal{C} \setminus T$, $h_{w_e}(x) = x$, then the leaf through x is finite since its intersection with D is the finite set $\bigcup_{v \in V(\mathcal{C})} h_{w_v}(x)$. This contradicts the fact that $x \in l_0$ since l_0 is infinite, and proves our claim.

Consider an oriented edge $e \in \mathcal{C} \setminus T$ and $x \in l_0$ such that $h_{w_e}([a, x]) \subsetneq [a, x]$. Since l_0 is singular, x is a subdivision point of $\mathcal{L}_i$ for i large enough. This does not necessarily prevent the existence of a combinatorial measure for $\mathcal{L}_i$ because a is not necessarily a subdivision point of $\mathcal{L}_i$. Up to replacing w_e by some power w of it, one can assume that the twisting morphism φ_w is trivial. Then for any integer m , there exists i large enough such that all segments $[x, h_w(x)], [h_w(x), h_w^2(x)], \dots, [h_w^{m-1}(x), h_w^m(x)]$ are mapped to each other under the holonomy of a power of w in $\mathcal{L}_i$. It follows that $\text{exponent}(\mathcal{L}_i)$ goes to infinity with i . □

7.1.3 Collapsing segments of measure 0, Rips band complex.

Now we assume that Σ is endowed with its topological foliation $\mathcal{F}$, and its non-atomic invariant transverse measure μ . Still, μ may fail to have full support. We now define a *Rips* band complex by collapsing connected components of the complement of the support of μ as follows.

Let D_μ be the quotient of D by the equivalence relation $\mu([x, y]) = 0$ (recall that μ has no atom), and $\pi_\mu : D \rightarrow D_\mu$ the quotient map. For each band B of Σ with bases J_0, J_1 , we consider the corresponding band B_μ with bases $\pi_\mu(J_0), \pi_\mu(J_1)$ and whose holonomy is induced by the holonomy of B . Let Σ_μ be the complex of bands on D_μ whose bands are the B_μ . The holonomy of the bands naturally defines a foliation $\mathcal{F}_\mu$ on Σ_μ . We still denote by μ the measure induced by μ on D_μ . This measure is invariant under the holonomy of $\mathcal{F}_\mu$.

Thus, one obtains a new foliated band complex Σ_μ , where one needs to generalise the definition of a band complex to allow some connected components of D_μ to be reduced to a point, and some bands to have bases reduced to a point. In this setting, we still can talk of leaves, singular leaves etc.

Since μ has no atom and has full support in D_μ , it can be transported to the Lebesgue measure of a finite union of intervals of $\mathbb{R}$ by a homeomorphism. Such a band complex Σ_μ with its measured foliation $(\mathcal{F}_\mu, \mu)$ is thus the object studied in [BF95] or in [GLP94] under the name system of isometries. We call such a band complex a *Rips* band complex.

We will use the same notations for the bands of Σ and the corresponding bands of Σ_μ . In particular, we will view any Σ -word as a Σ_μ -word, and conversely.

We now gather some simple facts about the projection $\Sigma \rightarrow \Sigma_\mu$.

Lemma 7.10. *Consider $x \in D_\mu$ whose $\mathcal{F}_\mu$ -leaf is singular (resp. pseudo-singular). Then there exists $\tilde{x} \in D$ with $\pi_\mu(\tilde{x}) = x$ such that $\tilde{x}$ lies in a singular (resp. pseudo-singular) leaf of $\mathcal{F}$.*

Proof. If x is a vertex of Σ_μ , then $\pi_\mu^{-1}(\{x\})$ contains a vertex of Σ and we are done. Otherwise, there is a Σ_μ -word w such that $[x, w)$ is a regular leaf path joining x to a vertex y of Σ_μ . Let $\tilde{y} \in \pi_\mu^{-1}(\{y\})$ be a vertex of Σ . Viewing w as a Σ -word, the point $h_{w^{-1}}(\tilde{y}) \in \pi_\mu^{-1}(\{x\})$ satisfies the lemma. If the leaf through x is pseudo-singular but not singular, let w be a Σ_μ word such that $h_w(x) = x$ and h_w reverses the orientation. Then the corresponding Σ -word w has a fix point $\tilde{x}$, so the leaf through $\tilde{x}$ is pseudo singular. Since x is the unique fix point of h_w , $\pi_\mu(\tilde{x}) = x$. $\square$

Lemma 7.11. *No regular leaf of Σ_μ is finite.*

Proof. Let $x \in D_\mu$ whose Σ_μ -leaf is regular and finite. Consider a small open interval $I \subset D_\mu$ containing x in its interior, small enough so that the every leaf meeting I is regular and finite. Let K be the union of leaves meeting I . Then $\pi_\mu^{-1}(K)$ is a union of regular leaves of Σ . Since μ is a limit of measures supported on singular leaves of Σ , $\mu(\pi_\mu^{-1}(I)) = 0$, a contradiction. $\square$

7.2 Moves

We describe moves that can be performed on a band complex together with a prelamination. These combinatorial moves can be performed by the algorithm. They can be performed in the forward direction on a prelaminated band complex, or in the backward direction on a naked band complex (without prelamination). Each of these forward moves has a counterpart which can be applied to a measured foliation. Performing a move on the foliation will give a way to uniformly perform a move on a sequence of prelaminations represented by this foliation.

Each band complex and prelamination comes with a set of rational constraints in standard form, described by $\rho : S_\pm^+ \rightarrow \mathcal{M}$ and a tuple $\underline{m} \in \mathcal{M}^{E(\mathcal{L})}$. We always assume that $\underline{m}$ is invariant under the holonomy and shows no incompatibility of rational constraints (see section 6.2.1). All the moves will keep ρ and $\mathcal{M}$ unchanged.

We introduce inert moves before restriction and extension moves (see Section 6.2.4 for definitions).




Figure 5: Moves

7.2.1 Band subdivision

Let Σ be a prelaminated band complex with domain D , $\mathcal{L}$ a prelamination on Σ , $B = [a, b] \times [0, 1]$ a band of Σ , and $\{c\} \times [0, 1] \subset B$ a leaf segment of $\mathcal{L}$ with $c \notin \{a, b\}$ (see Figure 5)

Let Σ' be the band complex with domain $D' = D$, obtained by replacing the band B by two bands $B'_a = [a, c] \times [0, 1]$ and $B'_b = [c, b] \times [0, 1]$, the attaching maps being the restrictions of the attaching maps of B . The twisting morphisms of B'_a and B'_b are the same as the twisting morphism of B . The prelamination $\mathcal{L}$ naturally induces a prelamination $\mathcal{L}'$ on Σ' . The set $\underline{m} \in \mathcal{M}^{E(\mathcal{L})}$ of rational constraints associated to $\mathcal{L}$ can be seen as a set of rational constraint on $\mathcal{L}'$ since $E(\mathcal{L}') = E(\mathcal{L})$. This induces a set of rational constraints on Σ' by writing each elementary segment of Σ' as a concatenation of elementary segments of $\mathcal{L}$.

It is clear that any solution of Σ inducing $\mathcal{L}$ defines a solution of Σ' inducing $\mathcal{L}'$, and that any solution of Σ' induces a solution of Σ . Thus, $(\Sigma, \mathcal{L}) \rightarrow (\Sigma', \mathcal{L}')$ is an inert move.

If we are given a measured foliation $(\mathcal{F}, \mu)$ on Σ instead of a prelamination, together with a leaf segment $\{c\} \times [0, 1] \subset B$, we can similarly subdivide the band to obtain a new measured foliation $(\mathcal{F}', \mu')$ on a new band complex Σ' . Moreover, if $\mathcal{F}$ represents an increasing sequence of prelaminations $\mathcal{L}_i$, and if $\{c\} \times [0, 1]$ is in a special leaf of $\mathcal{F}$ (as in Definition 7.1), then for all i large enough, $\{c\} \times [0, 1]$ is contained in a leaf segment of $\mathcal{L}_i$, so one can perform the band subdivision move on $(\Sigma, \mathcal{L}_i)$, and get an increasing sequence of prelaminations $\mathcal{L}'_i$ on Σ' represented in $\mathcal{F}'$.

Definition 7.12. Assume that the topological foliation $\mathcal{F}$ represents $\mathcal{L}_i$.

We say that a move $(\Sigma, \mathcal{F}) \rightarrow (\Sigma', \mathcal{F}')$ is compatible with $\mathcal{L}_i$ if for all i large enough, one can perform the corresponding combinatorial move $(\Sigma, \mathcal{L}_i) \rightarrow (\Sigma', \mathcal{L}'_i)$ (including rational constraints) and $\mathcal{L}'_i$ is represented by $\mathcal{F}'$.

We proved:

Lemma 7.13. Assume that $\mathcal{F}$ represents a sequence of prelaminations $\mathcal{L}_i$. If e is a leaf segment of $\mathcal{F}$ contained in a special leaf, then band subdivision along e is compatible with $\mathcal{L}_i$.

7.2.2 Band removal

Let $(\Sigma, \mathcal{L})$ be a prelaminated band complex with domain D . Assume that there is an oriented band $B = [a, b] \times [0, 1]$ with bases $J_0 = [a_0, b_0]$, $J_1 = [a_1, b_1]$, and a Σ -word w not involving the band B or its inverse, whose holonomy map h_w is defined and coincides with the holonomy of B on $\{a_0, b_0\}$ (see Figure 5).

Let $(\Sigma', \mathcal{L}')$ be the prelaminated band complex obtained by performing on $(\Sigma, \mathcal{L})$ a band subdivision for each leaf segment of $[a, w)$ and $[b, w)$, then by removing the band B and all the leaf segments contained in B . Clearly, any solution of Σ inducing $\mathcal{L}$ is a solution of Σ' inducing $\mathcal{L}'$.

If the twisting morphisms of w and B agree, then conversely, any solution of Σ' (without prelamination) is a solution of Σ . Consider the case where the twisting morphism $\varphi = \varphi_w^{-1} \varphi_B$ is non-trivial in $\text{Aut}(S_{\pm})$. Let $m_{J_0} \in \mathcal{M}$ be the element induced by the rational constraint $\underline{m}$ on J_0 . Since there is no incompatibility of rational

constraints, $m_{J_0} \in \rho(\text{Fix}\varphi)$. By definition of standard form for rational constraints, $\text{Fix}\varphi$ is represented by ρ so $\text{Fix}\varphi = \rho^{-1}(\rho(\text{Fix}\varphi))$. It follows that $\rho^{-1}(m_{J_0}) \subset \text{Fix}\varphi$ so the rational constraint m_{J_0} alone imposes the fact that any solution σ' of Σ' satisfies $\sigma_{J_0} \in \text{Fix}\varphi$. Thus, in this case too, any solution of Σ' is a solution of Σ .

If we are given a measured foliation $(\mathcal{F}, \mu)$ on Σ instead of a prelamination, and if h_w is defined and coincides with the holonomy of B on $\{a_0, b_0\}$ as above, we can similarly perform this move to obtain a new measured foliation $(\mathcal{F}', \mu')$ on a new band complex Σ' . If $\mathcal{F}$ represents an increasing sequence of prelaminations $\mathcal{L}_i$ without incompatibility of rational constraints, then a_0 and b_0 lie in singular, hence special leaves. Thus, for i large enough, the $\mathcal{L}_i$ -holonomy of w will be defined on a_0 and b_0 , and we will be able to perform the band removal move on $\mathcal{L}_i$. This defines an increasing sequence of prelaminations $\mathcal{L}'_i$ on Σ' represented in $\mathcal{F}'$. Thus, the move on the foliated band complex uniformly induces moves on the prelaminations. We thus proved:

Lemma 7.14. *Assume that $\mathcal{F}$ represents a sequence of prelaminations $\mathcal{L}_i$ without incompatibility of rational constraints. Assume that B is an oriented band and w a Σ -word not involving $B^{\pm 1}$ whose holonomy maps one base of B exactly to the other base of B .*

Then the band removal move on $\mathcal{F}$ is compatible with $\mathcal{L}_i$.

7.2.3 Domain cut

Let $(\Sigma, \mathcal{L})$ be a prelaminated band complex with domain D , and consider $x \in D \setminus \partial D$ a $\mathcal{L}$ -subdivision point, and assume that x is non-orphan. Let $B_1, \dots, B_k$ the oriented bands containing x in the interior of their domain (one may have $B_i = B_j^{-1}$ for some $i \neq j$). By performing a band subdivision of the band B_i along the leaf segment $[x, B_i)$, we can assume that x is in the interior of no base of band. Now cut D along x , i. e. replace D by D' the disjoint union of the closure of the connected components of $D \setminus \{x\}$. The attaching maps of the bands are still well defined, and $\mathcal{L}$ induces a prelamination $\mathcal{L}'$ on the obtained band complex Σ' .

It is clear that any solution of Σ inducing $\mathcal{L}$ defines a solution of Σ' inducing $\mathcal{L}'$, and that any solution of Σ' induces a solution of Σ . Thus, $(\Sigma, \mathcal{L}) \rightarrow (\Sigma', \mathcal{L}')$ is an inert move. It can also be viewed as a restriction move for the map $\iota : D' \rightarrow D$ induced by inclusion as long as x lies in the active part of D , the backwards move $\Sigma' \rightarrow \Sigma$ being an extension move of Lipschitz factor 1.

This move generalises naturally to a foliation $\mathcal{F}$ instead of a prelamination, and if x lies in a special leaf, then it is non-orphan in some $\mathcal{L}_i$, so it uniformly induces domain cut moves on a sequence of prelaminations represented by $\mathcal{F}$:

Lemma 7.15. *Assume that $\mathcal{F}$ represents a sequence of prelaminations $\mathcal{L}_i$.*

If $x \in D$ lies in a special leaf of $\mathcal{F}$, then domain cut at x is compatible with $\mathcal{L}_i$.

7.2.4 Pruning move

The pruning move for a prelaminated band complex has been described in section 6.2.4. This move generalises naturally to a foliation $\mathcal{F}$ instead of a prelamination, and this move is always compatible with a sequence of prelaminations represented by $\mathcal{F}$.

7.2.5 Forgetful move

Let $(\Sigma, \mathcal{L})$ be a prelaminated band complex with domain D . Assume that there is a connected component D_0 of D which contains exactly two bases of bands B_1, B_2 with $B_1 \neq B_2^{\pm 1}$, and such that $\text{dom } B_1 \cup \text{dom } B_2 = D_0$ and $I = \text{dom } B_1 \cap \text{dom } B_2$ is a non-degenerate segment.

By performing a pruning move on the components of $D_0 \setminus I$, we can assume that $\text{dom } B_1 = \text{dom } B_2 = D_0$. Let Σ' the band complex with domain $D \setminus D_0$, and where the bands B_1, B_2 are removed and replaced by a single band B' with holonomy $h_{B_2} \circ h_{B_1}^{-1}$ and twisting morphism $\varphi_{B_2} \circ \varphi_{B_1}^{-1}$. The prelamination $\mathcal{L}$ induces a prelamination $\mathcal{L}'$ on B' , whose set of subdivision points is $D_{\mathcal{L}} \cap D'$, and such that leaf segments of B' are exactly the segments $[x, B']$ such that $[x, B_2 B_1^{-1})$ is a leaf path of $\mathcal{L}$.

It is clear that any solution of Σ inducing $\mathcal{L}$ defines a solution of Σ' inducing $\mathcal{L}'$, and that any solution of Σ' induces a solution of Σ . Thus, for any partition $D = D_I \cap D_A$ where D_A contains both bases of B_1 and of B_2 , the move $(\Sigma, \mathcal{L}) \rightarrow (\Sigma', \mathcal{L}')$ is a restriction move with respect to the inclusion $\iota : D' \hookrightarrow D$ and $\Sigma' \rightarrow \Sigma$ is an extension move of Lipschitz factor 3.

This move generalises naturally to a foliation $\mathcal{F}$ instead of a prelamination, and it uniformly induces forgetful moves on a sequence of prelaminations represented by $\mathcal{F}$.

Lemma 7.16. *Assume that $\mathcal{F}$ represents a sequence of prelaminations $\mathcal{L}_i$. Consider D_0 a connected component of D which contains exactly two bases of bands B_1, B_2 with $B_1 \neq B_2^{\pm 1}$, and such that $\text{dom } B_1 \cup \text{dom } B_2 = D_0$, and $\text{dom } B_1 \cap \text{dom } B_2$ is a non-degenerate segment.*

Then the forgetful move on $\mathcal{F}$ (consisting in removing D_0) is compatible with $\mathcal{L}_i$.

7.3 Minimal components

A Rips band complex has a decomposition into a finite union of *minimal components* ([BF95, GLP94]). In this subsection, we introduce μ -minimal components of the topological foliation Σ which mimic the minimal components of the corresponding Rips band complex Σ_μ . We prove in this subsection an important result showing that the measure of an interval I controls the relative length of any solution inducing $\mathcal{L}_i$ for i sufficiently large.

Consider a foliation $\mathcal{F}$ on a band complex Σ with domain D . Recall that the singular leaf segments of $\mathcal{F}$ are the leaf segments in the boundary of the bands of Σ , and that a leaf path is regular if it does not involve any singular leaf segment. We say that two points are in the same $\mathcal{F}$ -leaf if they can be connected by a regular leaf path.

Thus, two points of Σ_μ are in the same $\mathring{\mathcal{F}}_\mu$ -leaf if they can be joined by a regular leaf path of $\mathcal{F}_\mu$ (but the projection in Σ_μ of a regular leaf path of $\mathcal{F}$ may fail to be regular in Σ_μ). It is convenient to call $\mu([x, y])$ the μ -distance between x and y .

Definition 7.17. *Let $(\Sigma_\mu, \mathcal{F}_\mu)$ be a Rips band complex such that no component of D and no base of band is reduced to one point.*

We say that $(\Sigma_\mu, \mathcal{F}_\mu)$ is minimal if for each $x \in D_\mu \setminus \partial D_\mu$, every $\mathring{\mathcal{F}}_\mu$ -leaf is dense in Σ_μ .

Definition 7.18. A measured foliation $(\mathcal{F}, \mu)$ on a band complex Σ with domain D is called μ -minimal if all connected components of D and all bands of Σ have positive measure, and

- Σ_μ is minimal (as a Rips band complex)
- for each $x \in \partial D$, there exists a Σ -word whose holonomy is defined on an interval of positive measure, and maps x at positive μ -distance of ∂D .

Remark 7.19. One easily checks that all leaves of a μ -minimal band complex are infinite, but this may be false if one removes the last condition of the definition. Leaves of a μ -minimal band complex may in general fail to be dense, as there may be wandering intervals.

Assume that Σ is a μ -minimal foliated band complex representing a sequence of prelaminations $\mathcal{L}_i$. Given a solution σ of Σ inducing $\mathcal{L}_i$ for i large enough, one can control the ratio $|\sigma_I|/|\sigma|$ in terms of the measure $\mu(I)$:

Lemma 7.20. Let $(\Sigma, \mathcal{F}, \mu)$ be a μ -minimal foliated band complex with domain D . Let $\mathcal{L}_i$ be a sequence of prelaminations represented by $\mathcal{F}$.

For all $\varepsilon > 0$, there exists $\eta > 0$ such that the following holds: for each segment $I \subset D$ with $\mu(I) < \eta$, and for i large enough, there exists $I' \supset I$ adapted to $\mathcal{L}_i$, and a certificate of ε -shortness of I' relative to D as in Definition 6.11 (ensuring that $\frac{|\sigma_{I'}|}{|\sigma|} \leq \varepsilon$ by Lemma 6.12).

Proof. Fix $\varepsilon > 0$, and $N \geq \frac{1}{\varepsilon}$. For $x \in D_\mu \setminus \partial D_\mu$, the $\mathring{\mathcal{F}}_\mu$ -leaf of x is infinite, so we can find N distinct Σ_μ -words $w_1, \dots, w_N$ whose holonomies are defined on a neighbourhood of x , and map x to N distinct points of D_μ . Take $\varepsilon_1, \varepsilon_2 > 0$ small enough so that the holonomies h_{w_i} are defined on $[x - \varepsilon_1, x + \varepsilon_2]$ and map this segment to N disjoint intervals. Since every leaf is dense in Σ_μ , one can assume moreover that $x - \varepsilon_1$ and $x + \varepsilon_2$ lie in singular leaves of Σ_μ . Let $I_x \subset D$ whose endpoints are in singular leaves of Σ , and such that $\pi_\mu(I_x) = [x - \varepsilon_1, x + \varepsilon_2]$. Then for i large enough, I_x is adapted to $\mathcal{L}_i$, and has a certificate of ε -shortness relative to D .

Consider $x \in \partial D_\mu$ and $\tilde{x}$ its preimage in ∂D . Consider w a Σ -word whose $\mathcal{F}$ -holonomy is defined on an interval of positive measure, and maps $\tilde{x}$ at positive μ -distance from ∂D . Let $\tilde{y} = h_w(\tilde{x})$, and $y = h_w(x)$ the image of $\tilde{y}$ in D_μ . Consider the interval $I_y \subset D$ defined above and define $I_x = h_w^{-1}(I_y)$. The endpoints of I_x are in singular leaves of Σ , so I_x is adapted to $\mathcal{L}_i$ for i large enough, and has a certificate of ε -shortness relative to D .

Consider the covering of D_μ by the interiors in D_μ of $\pi_\mu(I_x)$, and by compactness, consider η the Lebesgue number of this covering. Then any segment $J \subset D_k$ of measure at most η is contained in the interior of some I_x , and the lemma follows. $\square$

Proposition 7.21. Consider a band complex Σ with a measured foliation $(\mathcal{F}, \mu)$ without atom representing a sequence of prelamination $\mathcal{L}_i$. Assume that there is a bound on the exponent of periodicity of $\mathcal{L}_i$.

Then one can perform a finite number of inert moves on $(\Sigma, \mathcal{F})$ compatible with $\mathcal{L}_i$ so that the obtained foliated complex $(\Sigma', \mathcal{F}')$ is the disjoint union of finitely many μ -minimal band complexes $\Lambda_1, \dots, \Lambda_p$ together with finitely many bands and components of D' of measure 0.

The exponent of periodicity of $\mathcal{L}_i$ is defined in Definition 6.5.

Proof. We first recall the decomposition into minimal components for Σ_μ . Let $\mathcal{C}_\mu \subset \Sigma_\mu$ be the finite graph defined by the union of singular finite $\mathring{\mathcal{F}}_\mu$ -leaves (a $\mathring{\mathcal{F}}_\mu$ -leaf is *singular* if it contains a subdivision point of Σ_μ , or equivalently, if the $\mathcal{F}_\mu$ -leaf containing it is singular). This graph is finite because Σ_μ has only finitely many vertices.

Denote by $\Lambda_{\mu,1}, \dots, \Lambda_{\mu,p}$ the Rips band complexes obtained as the closure of the connected components of $\Sigma_\mu \setminus \mathcal{C}_\mu$. Equivalently, $\Lambda_{\mu,i}$ are the connected components of the band complex obtained from Σ_μ by performing band subdivision and domain cuts at edges and vertices of $\mathcal{C}_\mu$, and by removing bands of width 0. By [GLP94, BF95], either $\Lambda_{\mu,i}$ is minimal, or $\Lambda_{\mu,i}$ is a *family of finite leaves* (all leaves of $\Lambda_{\mu,i}$ are finite in particular). In our setting, families of finite leaves cannot occur by Lemma 7.11.

We now construct a lift $\mathcal{C}$ of $\mathcal{C}_\mu$ in Σ . Working independently on each connected component on $\mathcal{C}_\mu$, we can assume that $\mathcal{C}_\mu$ is connected. Consider $x_\mu \in \mathcal{C}_\mu$, and $\tilde{x} \in D$ be a point whose leaf is special with $\pi_\mu(\tilde{x}) = x_\mu$. Existence of $\tilde{x}$ follows from Lemma 7.10 since singular leaves are special. Consider T a maximal tree of $\mathcal{C}_\mu$. Given two vertices $v_1, v_2 \in T$, we denote by $[x_1, x_2]_T$ the path of T joining them. First, we can lift T in Σ . Each segment $[x_\mu, v]_T$ defines a Σ -word w_v , and $\tilde{x}$ lies in the interior of $\text{dom } w_v$ since $\mathcal{C}_\mu$ contains no singular leaf segment. Therefore, one can define $\tilde{T} \subset \Sigma$ as the union of the leaf paths $(\tilde{x}, w_v)$. For each oriented edge $e \subset \mathcal{C}_\mu \setminus T$ joining v_1 to v_2 , consider B_e the oriented band containing it, and the Σ -word $w_e = w_{v_2}^{-1} B_e w_{v_1}$. To lift e , we need that the endpoint of the leaf segment $(\tilde{v}_1, B_e)$ coincides with $\tilde{v}_2$, where $\tilde{v}_i$ denotes the lift of v_i in $\tilde{T}$. Equivalently, we need to check that for all edge e of $\mathcal{C}_\mu \setminus T$, $h_{w_e}(\tilde{x}) = \tilde{x}$.

Let $I \subset D$ be the preimage of x_μ under π_μ . Since $\tilde{x}$ and $h_{w_e}(\tilde{x}) \in I$, we are done if I is reduced to one point. As the holonomy in Σ_μ of every loop in $\mathcal{C}_\mu$ based at x_μ is defined on a neighbourhood of x_μ , this defines in Σ a morphism $\psi : \pi_1(\mathcal{C}, x_\mu) \rightarrow \text{Homeo}(I)$. Let H be the image of ψ and $H_+ < H$ the orientation preserving subgroup.

We claim that for each point $\tilde{y} \in I$ whose leaf is special, $h.\tilde{y} = \tilde{y}$ for all $h \in H_+$. Otherwise, one can assume for instance $h.\tilde{y} > \tilde{y}$, so $h^n.\tilde{y} > \tilde{y}$ for all $n > 0$. Moreover, up to changing h to a power, we may assume that $h = \psi(w)$ for some Σ -word w with trivial twisting morphism. Since $\tilde{y}$ lies in a special leaf of $\mathcal{F}$, for any $n > 0$ there exists i such that the all the intervals $[\tilde{y}, h.\tilde{y}], \dots, [h^{n-1}.\tilde{y}, h^n.\tilde{y}]$ are equivalent under the holonomy of $\mathcal{L}_i$, so $\text{Exponent}(\mathcal{L}_i) \rightarrow \infty$, a contradiction.

Let $w \in \pi_1(\mathcal{C}, x_\mu)$ be a Σ -word whose holonomy does not fix $\tilde{x}$. Since $\tilde{x}$ lies in a singular hence special leaf of $\mathcal{F}$, the homeomorphism $h = \psi(w)$ reverses the orientation. Let $m \in I$ be its unique fixed point. The leaf through m is special since w and $[x_\mu, h_w(x_\mu)]$ define a Möbius strip in some $\mathcal{L}_i$, whose core needs to appear in some $\mathcal{L}_j$. The claim above shows that $H_+.m = m$, and since $H = \langle H_+, h_w \rangle$, $H.m = m$. This means that choosing m instead of $\tilde{x}$ as a lift of x_μ , one can lift $\mathcal{C}_\mu$ to Σ .

Let $\mathcal{C} \subset \Sigma$ be the graph obtained by lifting each connected component of $\mathcal{C}_\mu$ as above. Let $(\Sigma', \mathcal{F}')$ be the measured foliated band complex with domain D' obtained from $(\Sigma, \mathcal{F})$ by first performing a band subdivision at each edge of $\mathcal{C}$ and then a domain cut at each vertex of $\mathcal{C}$. We still denote by μ the obtained transverse measure on Σ' . The band subdivisions along edges of $\mathcal{C}$ are mirrored by band subdivisions on Σ_μ since edges of $\mathcal{C}_\mu$ are at positive μ -distance of the boundary of the band containing

them. This is not true for the domain cuts: given a vertex $x \in \mathcal{C}$, and an oriented band B containing x in the interior of domain, the leaf segment $[x, B)$ may fail to lie in $\mathcal{C}$, in which case x is at zero μ -distance from an endpoint of $\text{dom } B$; in this case, the domain cut at x will split the band B into two bands, one of them having measure 0. Thus, Σ'_μ differs from Σ_μ only by the potential presence of additional bands of measure 0. Let D'^0 be the union of connected components of D' with positive measure, and Σ'^0 the band complex on D'^0 whose bands are the bands of Σ' with positive measure. Then Σ'^0 is a disjoint union of foliated band complexes $\Lambda'_1, \dots, \Lambda'_p$, such that $\Lambda'_{1,\mu}, \Lambda'_{p,\mu}$ are precisely the minimal components of Σ_μ .

We need to perform some more operations on Σ' to make sure that Λ'_i satisfies the second condition of μ -minimality. We focus on a component $\Lambda'_i \subset \Sigma'$, and to make notations lighter, we use the notations $\Lambda = \Lambda'_i$, and by $D \subset D'$ its domain. Consider the graph $\mathcal{D}_\mu \subset \Lambda_\mu$ with vertex set ∂D_μ , and whose edges are the leaf segments of Λ_μ having both endpoints in ∂D_μ . Working independently on each component of $\mathcal{D}_\mu$, we can assume that $\mathcal{D}_\mu$ is connected. There exists a vertex $x_\mu \in \mathcal{D}_\mu$ and an oriented band B_μ of Σ_μ sending x_μ into the interior of D_μ : otherwise leaves through the points close to the vertex set of $\mathcal{C}_\mu$ would be compact.

Let $\tilde{x}$ be the (unique) endpoint of D projecting to x_μ . A loop in $\mathcal{D}_\mu$ based at x_μ defines a Σ -word w whose holonomy h_w in Σ is defined on an interval of positive measure, and contains a point at μ -distance 0 from $\tilde{x}$. Moreover, h_w preserves the orientation. We claim that for each $y \in \text{dom } w$ and in a special leaf, $h_w(y) = y$. Indeed, write $\text{dom } w = [a, b]$, and assume for instance $h_w(y) > y$. If $\mu[y, b] > 0$, then $h_{w^k}(y)$ is defined for all $k > 0$ since $\mu([y, h_w(y)]) = 0$; if $\mu[y, b] = 0$, then up to changing w to w^{-1} and y to $h_w(y)$, the same conclusion holds. This contradicts the bound on the exponent of periodicity as above.

We construct a lift $\mathcal{D}$ of $\mathcal{D}_\mu$. Let T be a maximal tree of $\mathcal{D}_\mu$. For each $v \in \mathcal{D}_\mu$, let w_v be the Σ -word representing the segment $[x_\mu, v]_T$. For each oriented edge $e \in \mathcal{D}_\mu \setminus T$ joining v_1 to v_2 , let B_e be the oriented band containing it, and $w_e = w_{v_2}^{-1} B_e w_{v_1}$. Let $I \subset D'^0$ be the intersection of all domains $\text{dom } w_v$, $\text{dom } w_e$, and $\text{dom } B$. Since all these domains have positive measure and a point a μ -distance 0 from $\tilde{x}$, I has an endpoint $\tilde{y} \in \pi_\mu^{-1}(x_\mu)$. The leaf through $\tilde{y}$ is singular hence special, so $h_{w_e}(\tilde{y}) = \tilde{y}$ for all e by the claim above. Thus, the union of all leaf paths $[\tilde{y}, w_e)$, $[\tilde{y}, w_v)$ defines a lift $\mathcal{D}$ of $\mathcal{D}_\mu$.

Perform on Σ band subdivisions at all edges of $\mathcal{D}$ and domain cuts at all vertices of all connected components of $\mathcal{D}$, and denote by Σ'' with domain D'' the obtained foliated band complex. Let D''^0 be the union of connected components of D'' with positive measure, and Σ''^0 the band complex on D''^0 whose bands are the bands of Σ'' with positive measure. Each band of Σ' has been cut into one band in Σ''^0 , and at most two bands of measure 0 (at most one on each side). A similar fact holds for connected components of D' . Thus Λ has been transformed to $\Lambda'' \subset \Sigma''^0$ together with some intervals and bands of measure 0.

We claim that Λ'' is a μ -minimal component. Denote by $D_{\Lambda''} \subset D''^0$ the domain of Λ'' . Because of the cutting operation performed, for every endpoint x of a base of a band of Λ'' , either x lies in $\partial D_{\Lambda''}$, or it lies at positive μ -distance from it. Since for every endpoint x_μ of D_μ there is a Σ_μ -word defined on a set of positive measure mapping x_μ to the interior of D_μ , this word lifts to a Λ'' -word mapping the corresponding endpoint of $D_{\Lambda''}$ at positive μ -distance from $\partial D_{\Lambda''}$.

Since all band subdivision moves and all domain cuts have been performed on special leaves of $\mathcal{F}$, these moves are compatible with $\mathcal{L}_i$, and the proposition follows.

□

7.4 Homogeneous components

Recall that two points of Σ_μ are in the same $\mathring{\mathcal{F}}_\mu$ -leaf if they can be joined by a regular leaf path of $\mathcal{F}_\mu$.

Definition 7.22. *One says that Σ_μ contains a homogeneous component (also called axial or toral component) if there exists a non-degenerate open interval $I \subset D_\mu$ and a subgroup $P \subset \text{Isom}(\mathbb{R})$ with dense orbits such that for all $x, y \in I$, x, y are in the same $\mathring{\mathcal{F}}_\mu$ -leaf if and only if x, y are in the same P -orbit.*

When a minimal component satisfies this definition, it is called a homogeneous component. If Σ_μ contains a homogeneous component as in the definition above, one of its minimal components is a homogeneous component.

Proposition 7.23. *If Σ_μ contains a homogeneous component, then the exponent of periodicity of $\mathcal{L}_i$ goes to infinity as i goes to infinity.*

Given a Σ_μ -word w whose holonomy is a translation of positive length $t(w)$, whose domain of definition has positive measure, its *translation ratio* is the ratio $tr(w) = \mu(\text{dom}(w))/t(w)$. The proposition is based on the following fact from Rips' Theory (of which we give a proof below):

Fact. *Assume that a system of isometries Σ_μ contains a homogeneous component. Then there are Σ_μ -words with arbitrarily large translation ratio.*

Proof of Proposition 7.23. Let k be the cardinal of $\text{Aut}(S_\pm)$. Since $tr(w^k) \geq \frac{1}{k}tr(w) - 1$, there are Σ_μ -words with trivial twisting whose translation ratio is arbitrarily large. Consider such a Σ_μ -word w with translation ratio at least N , and view it as a Σ -word. Let $x \in D$ be an endpoint of $\text{dom } w$. Since $tr(w) \geq N$, up to changing w to w^{-1} , we can assume that for all $k = 0, \dots, N-1$, $x \in \text{dom } w^k$. Since the leaf through x is singular hence special, for all i large enough, all intervals $[x, h_w(x)], \dots, [h_w^{N-2}(x), h_w^{N-1}(x)]$ are equivalent under the holonomy of some w^j . The exponent of periodicity of $\mathcal{L}_i$ then goes to infinity with i . □

Proof of the fact. Given $\eta > 0$, consider Σ' the Rips band complex on D_μ obtained by narrowing each band by η on each side. More formally, this amounts to restricting $f_{B,\varepsilon} : [a, b] \times \{\varepsilon\} \rightarrow D$ to $[a+\eta, b-\eta] \times \{\varepsilon\}$. If Σ_μ contains a homogeneous component, then one can find $\eta > 0$ such that Σ' has an infinite orbit ([GLP94]). In particular, there exists Σ' -words whose holonomy is a translation of arbitrarily small length. Fix some large $M \in \mathbb{R}$, and consider w whose holonomy is a translation of length at most η/M . The holonomy of corresponding Σ_μ -word is a translation of the same length, and its domain has measure at least 2η , so $tr(w) \geq 2M$. □

7.5 Independent bands

A theorem by Gaboriau asserts that if a Rips band complex Σ_μ has no homogeneous component, one can narrow its bands so that its bands are *independent*, meaning that no $\mathring{\mathcal{F}}_\mu$ -leaf contains a cycle. This result was also used in [GLP94] or [Gui98, Section 6] for the analysis of actions on $\mathbb{R}$ -trees. Our goal is to mirror this fact in the topological foliated band complex $(\Sigma, \mathcal{F})$, and into the prelaminated band

complexes $(\Sigma, \mathcal{L}_i)$.

Given a foliated band complex $(\Sigma, \mathcal{F})$ with domain D , an *orbit* of $(\Sigma, \mathcal{F})$ is the intersection of a leaf with D .

Theorem 7.24 ([Gab97]). *Given Σ_μ a Rips band complex with domain D without homogeneous component, there exists a Rips band complex $\Sigma'_\mu \subset \Sigma_\mu$ with same domain D , obtained by replacing each band $B = [a, b] \times [0, 1]$ of Σ by a narrower band $B' = [a', b'] \times [0, 1] \subset B$, and such that*

1. Σ_μ and Σ'_μ have the same orbits
2. Σ'_μ has independent bands: no $\mathring{\mathcal{F}}'_\mu$ -leaf contains a cycle
3. any singular leaf of Σ'_μ is contained in a pseudo-singular of Σ_μ .

Remark 7.25. If bands of Σ'_μ are independent, every pseudo-singular leaf is singular.

The last condition is not stated in Gaboriau's theorem but can be ensured by slightly modifying Gaboriau's construction.

Proof. We explain how to ensure that the set of pseudo-singular orbits of Σ_μ does not increase during Gaboriau's construction in the case where Σ_μ is a minimal component. For convenience, we give references to statements in Gaboriau's paper. To lighten notations, we drop the index μ so we denote by Σ the initial Rips band complex. Clearly, one can ensure that condition 3 holds if all orbits are finite ([Gab97, Lemme 5.2(1)]). The proof of Gaboriau's theorem (see [Gab97, section 7]) consists in modifying the band complex Σ by narrowing then re-enlarging bands. Instead of narrowing bands by a uniform amount, we need to choose narrowing widths so that we don't create new pseudo-singular leaves.

The set of sides of bands (i. e. the set of singular leaf segments) is the set of edges $E(\Delta)$ of the finite graph Δ of [Gab97]. Given a tuple $\underline{t} = (t_\varepsilon)_{\varepsilon \in E(\Delta)}$ of (small enough) non-negative numbers, one can define $\Sigma_{\underline{t}}$ by narrowing each band by t_ε on the side corresponding to ε .

Consider $\mathcal{T}$ a maximal subforest of Δ , a small tuple $\underline{t}$ so that $t_\varepsilon = 0$ for $\varepsilon \in \mathcal{T}$, and for $\varepsilon \notin \mathcal{T}$, choose $t_\varepsilon > 0$ so that the boundary of the new band lies in a singular leaf of Σ . Moreover, $\underline{t}$ should be small enough so that Σ and $\Sigma_{\underline{t}}$ have the same orbits ([Gab97, Prop. 4.5]). This is possible because the union of singular leaves is dense.

Let t_1 be the minimum of t_ε for $\varepsilon \notin \mathcal{T}$, and consider $\underline{u} = (u_\varepsilon)_{\varepsilon \in E(\Delta)}$ a small tuple with $u_\varepsilon = 0$ for $\varepsilon \notin \mathcal{T}$ and $0 < u_\varepsilon \leq t_1$ for $\varepsilon \in \mathcal{T}$ so that the singular leaves of $\Sigma_2 = \Sigma_{\underline{u} + \underline{t}}$ are contained in singular leaves of Σ . Since Σ is non-homogeneous, and since all coordinates of $\underline{u} + \underline{t}$ are positive, leaves of $\Sigma_{\underline{u} + \underline{t}}$ are compact ([Gab97, Th 3.3]). Denote by $e(\Sigma)$ the measure of the space of leaves of a Rips band complex Σ (this is 0 if every leaf is dense), and by $l(\Sigma)$ the sum of the transverse measure of its bands. Then $e(\Sigma_{\underline{t}}) = 0$, and using [Gus00, Th II.E.4], $e(\Sigma_{\underline{u} + \underline{t}}) = |\underline{u}|$ where $|\underline{u}| = \sum_{\varepsilon \in \Delta} u_\varepsilon$ (this is the generalisation we need of [Gab97, Prop. 4.3]). Since Σ_2 has compact leaves, one can find $\underline{v}$ such that $v_\varepsilon = 0$ for $\varepsilon \in \mathcal{T}$, and $\Sigma_3 = \Sigma_{\underline{u} + \underline{t} + \underline{v}}$ has the same orbits as Σ_2 , and has independent bands and has no new pseudo-singular leaves [Gab97, Lemme 5.2]. Since bands are independent, $e(\Sigma_3) + l(\Sigma_3) = \mu(D)$ ([Gab97, Th. 6.3]).

Finally, consider $\Sigma_4 = \Sigma_{\underline{t} + \underline{v}}$, obtained from Σ_3 by enlarging the bands by $\underline{u}$. Then as in [Gab97, section 7], Σ_4 has the same orbits as Σ , so $e(\Sigma_4) = 0$. Since $l(\Sigma_4) + e(\Sigma_4) = l(\Sigma_3) + |\underline{u}| = \mu(D)$, its bands are independent [Gab97, Th. 6.3].

Finally, one easily checks that any singular leaf of Σ_4 is contained in a pseudo-singular leaf of Σ . $\square$

Definition 7.26. *Say that a measured foliation $(\mathcal{F}, \mu)$ on Σ has μ -independent bands if the holonomy of no non-trivial reduced Σ_μ -word fixes an interval of positive measure.*

Our goal is the following proposition.

Proposition 7.27. *Consider a band complex Σ with a measured foliation $(\mathcal{F}, \mu)$ without atom representing a sequence of prelamination $\mathcal{L}_i$. Assume that there is a bound on the exponent of periodicity of $\mathcal{L}_i$, and that each $\mathcal{L}_i$ has an invariant combinatorial measure.*

Then one can perform a finite number of inert moves on $(\Sigma, \mathcal{F})$ compatible with $\mathcal{L}_i$ so that the obtained foliated complex $(\Sigma', \mathcal{F}')$ has μ -independent bands.

We will use the following useful finiteness property:

Proposition 7.28 (Segment-closed property, [GLP95]). *Let $(\Sigma_\mu, \mathcal{F}, \mu)$ be a Rips band complex, and $\varphi : I \rightarrow J$ a partial isometry between two segments of D_μ such that for all $x \in I$, x and $\varphi(x)$ are in the same leaf.*

Then there exists finitely many Σ_μ -words $w_1, \dots, w_n$ whose domains cover I , and whose holonomies coincide with φ on I . $\square$

Proof of Proposition 7.27. For each band B of Σ , we denote by B_μ the corresponding band of Σ_μ . Consider Σ'_μ the Rips band complex on D_μ whose bands $B'_\mu \subset B_\mu$ are given by Theorem 7.24. Consider a band B and I, J (resp. I_μ, J_μ and I'_μ, J'_μ) the bases of B (resp. B_μ, B'_μ). Let K_μ be the closure of a connected component of $I_\mu \setminus I'_\mu$. Since Σ_μ and Σ'_μ have the same orbits, the holonomy h_{B_μ} of B_μ is a partial isometry such that x and $h_{B_\mu}(x)$ are in the same Σ'_μ -leaf. By segment-closed property, there exist Σ'_μ -words $w'_1, \dots, w'_n$ whose domains cover K_μ , and whose holonomies coincide with h_{B_μ} on K_μ . We can assume that $\text{dom } w'_i$ is not reduced to a point i. e. has positive measure.

We now lift this situation to Σ . For each band B of Σ whose bases have positive measure, consider a restriction B' of B in Σ whose base projects to B'_μ in Σ_μ . By Lemma 7.10, one can choose the endpoints of B' in pseudo-singular leaves of $\mathcal{F}$. By remark 7.2, these leaves are special. Using the band subdivision move, one can subdivide the band B along the boundary of the corresponding sub-band B' . Doing this for each band of Σ gives get a new band complex Σ_1 whose set of bands is $\mathcal{B}' \sqcup \mathcal{B}''$ where $\mathcal{B}'$ is the set chosen lifts of bands of Σ'_μ .

For each band $B'' \in \mathcal{B}''$, there are Σ'_μ -words $w'_1, \dots, w'_n$ whose domains cover $\text{dom } B''_\mu$, and whose holonomies coincide with $h_{B''_\mu}$. We view these words as Σ_1 -words involving only bands of $\mathcal{B}'$. The domains of w'_i cover all $\text{dom } B''$ except for some finite union of intervals of measure zero. By further subdividing B'' along some singular leaves (defined as the boundary of the domain of some w'_i), we can replace each band B'' by a finite set of smaller sub-bands B''' so that either $\mu(\text{dom } B''') = 0$ or $\text{dom } B''' \subset \text{dom } w'_i$ for some i .

Fix some band B''' whose domain $[a, b]$ has positive measure. By construction, the holonomy of the word $u = w_i^{-1} B'''$ induces the identity on $\pi_\mu([a, b])$. If $h_u(a) \neq a$, then up to changing u to u^{-1} , $h_{u^k}(a)$ is defined for all $k > 0$. Since the leaf through a is special, this clearly implies that $\text{exponent}(\mathcal{L}_i)$ goes to infinity. It follows that h_u

fixes both a and b , so one can remove the band B''' using the band removal move. The bands of the obtained foliated band complex Σ_2 are either bands of $\mathcal{B}'$, or bands whose bases have measure zero. Thus, Σ_2 has μ -independent bands. $\square$

7.6 The pruning process

Consider a foliated band complex $(\Sigma, \mathcal{F}, \mu)$ with μ -independent bands, decomposed into μ -minimal components as in Proposition 7.21. Let Λ be a minimal component of this decomposition.

We recall the following pruning process (also known as process 1 of the Rips machine). All the moves are performed on Σ , modifying only $\Lambda \subset \Sigma$.

We denote by $D_\Lambda = D \cap \Lambda$ the domain of Λ , which will be the *active* part for the moves. The pruning process consists in the iteration of the following steps:

- *Step a.* Say that an open interval $I \subset D_\Lambda$ is *prunable* if every $x \in \overset{\circ}{I}$ lies in exactly one base of a band of Σ . If Λ_μ has no prunable interval of positive measure, the process stops. Otherwise, let $I_1, \dots, I_n$ be the complete list of maximal prunable intervals of positive measure, and perform a pruning move for each I_i .
- *Step b.* If some component of D_Λ contains at most one base of band, prune this band, and iterate step b as many times as possible.
- *Step c.* Perform on all possible forgetful moves on Λ .

Remark 7.29. In step a, note that each band of Σ having a base in D_Λ either lies in Λ or has measure zero. Step b and step c can be repeated only finitely many times in a row since they decrease the number of bands.

Starting with $(\Sigma_0, \mathcal{F}_0, \mu_0) = (\Sigma, \mathcal{F}, \mu)$, we define inductively $(\Sigma_{j+1}, \mathcal{F}_{j+1}, \mu_{j+1})$ by performing the three steps a,b,c on $(\Sigma_j, \mathcal{F}_j, \mu_j)$. The sequence of prelaminations $\mathcal{L}_i \subset \Sigma$ represented by $\mathcal{F}$ induces a sequence of prelaminations $\mathcal{L}_{j,i} \subset \Sigma_j$ represented in $\mathcal{F}_i$. Moreover, the rational constraints on $\mathcal{L}_i$ (in standard form) induce rational constraints on $\mathcal{L}_{j,i}$ which define rational constraints $\underline{m}_j \in \mathcal{M}^{E(\Sigma_j)}$ on Σ_j independently of i by compatibility of the rational constraints. All rational constraints are in standard form, using the same morphism $\rho : S_\pm^+ \rightarrow \mathcal{M}$.

We denote by $\Lambda_j \subset \Sigma_j$ the subset corresponding to the minimal component under study. We denote by $D_{\Sigma_j} \subset D$ and $D_{\Lambda_j} = D_{\Sigma_j} \cap \Lambda$ the domains of Σ_j and Λ_j , viewed as subsets of D .

Since bands are independent (which excludes homogeneous components), surface and exotic components can be defined by the fact that the pruning process stops or not ([GLP94]):

Definition 7.30. *Say that Λ is exotic (or Levitt, or thin), if the pruning process never stops. Otherwise it is a surface component.*

7.6.1 Exotic components

We focus on an exotic minimal component Λ of this decomposition. The goal of this section is the following proposition:

Proposition 7.31. *Consider a foliated band complex $(\Sigma, \mathcal{F}, \mu)$ with μ -independent bands, decomposed into μ -minimal components as in Proposition 7.21. Consider Λ an exotic μ -minimal component of Σ . Let $\mathcal{L}_i$ be a sequence of prelaminations represented by $\mathcal{F}$.*

Then for i large enough, there exists a shortening sequence of moves for $(\Sigma, \mathcal{L}_i)$ as in definition 6.13, certifying that a shortest solution of Σ cannot induce $\mathcal{L}_i$.

Proposition 7.31 will follow from the two following facts.

Lemma 7.32. *The complexity of Σ_j , defined as the sum of number of bands of Σ_j and of the number of connected components of D_j remains bounded.*

In particular, only finitely many distinct unfoliated band complexes with rational constraints appear in the sequence $\Sigma_1, \Sigma_2, \dots$.

Proof. The second assertion easily follows from the first one since all rational constraints are in standard form, defined using the same morphism $\rho : S_{\pm}^+ \rightarrow \mathcal{M}$.

We need only to prove that the number of bands of Λ_j and of connected components of D_{Λ_j} is bounded. The band complex Λ_j is homotopy equivalent to a graph Γ_j whose set of vertices $V(\Gamma_j)$ is the set of connected components of D_{Λ_j} and whose set of non-oriented edges $E(\Gamma_j)$ corresponds to non-oriented bands. The first Betti number of Λ_j is $b_1(\Lambda_j) = \#V(\Gamma_j) - \#E(\Gamma_j)$. The moves of the pruning process preserve the first Betti number: this is clear for moves involved in step b and c; for moves in step a, this follows from the fact that if $[a, b]$ is pruned, then either $a \in \partial D$, or a lies in the interior of the pruned band because otherwise, the $\mathring{\mathcal{F}}$ -leaf through a would be finite, contradicting μ -minimality.

After step b, every terminal vertex of Γ_j corresponds to a connected component of D_{Λ_j} containing a base of a band of $\Sigma_j \setminus \Lambda_j$. Similarly, after step c, every vertex of valence 2 of Γ_j corresponds to a connected component of D_{Λ_j} containing a base of a band of $\Sigma_j \setminus \Lambda_j$. Since step c does not create new terminal vertices, the number of vertices of valence 1 and 2 of Γ_j is bounded. Since $b_1(\Gamma_j)$ is bounded, the lemma follows. $\square$

Lemma 7.33. *For all $\eta > 0$, there exists j large enough such that $\mu(D_{\Lambda_j}) < \eta$.*

Proof. By [GLP94, Prop. 7.1], the μ -diameter of the connected components of D_{Λ_j} converges to 0. Since the number of connected components of D_{Λ_j} is bounded, the proposition follows. $\square$

Proof of Proposition 7.31. Since the complexity of band complexes appearing in the sequence $\Sigma_1, \Sigma_2, \dots$ is bounded, there is a bound C on the number of unfoliated band complexes with rational constraints appearing in this sequence. The number of moves performed at step a, b and c, is bounded in terms of the complexity of Σ_j , and is therefore bounded by some number k . It follows that for all j , one can go back from the unfoliated band complex Σ_j to the initial unfoliated band complex Σ using at most Ck extension moves.

Consider $\varepsilon < 1/3^{Ck}$. Using Lemma 7.20, consider η so that for any $I \subset D_{\Lambda}$ with $\mu(I) < \eta$, there exists ε -shortness certificates for I relative to D_{Λ} . By Lemma 7.33, let j_0 be large enough so that $\mu(D_{\Lambda_{j_0}}) < \eta$.

Let $\mathcal{L}_i$ be a sequence of prelaminations represented in $\mathcal{F}$, and consider i_0 large enough so that the sequence of moves from $(\Sigma, \mathcal{F})$ to $(\Sigma_{j_0}, \mathcal{F}_{j_0})$ can be applied to $(\Sigma, \mathcal{L}_{i_0})$, transforming it into $(\Sigma_{j_0}, \mathcal{L}')$. These moves are restriction moves with inert

part $D_I = D \setminus \Lambda$ and active part $D_A = D_\Lambda$. Since Σ can be obtained from Σ_{j_0} using at most Ck extension moves of Lipschitz factor 3, going from Σ_{j_0} to Σ is an extension move of Lipschitz factor 3^{Ck} . Thus, we have found a shortening sequence of moves for $(\Sigma, \mathcal{L}_{i_0})$, which certifies that a shortest solution of Σ cannot induce $\mathcal{L}_{i_0}$ (see Definition 6.13). $\square$

7.6.2 Surface components

Proposition 7.34. *Consider a foliated band complex $(\sigma, \mathcal{F}, \mu)$ with μ -independent bands, decomposed into μ -minimal components as in Proposition 7.21. Assume that Λ is a μ -minimal surface component. Let $\mathcal{L}_i$ be a sequence of prelaminations represented by $\mathcal{F}$.*

Then for i large enough, there exists a shortening sequence of moves for $(\Sigma, \mathcal{L}_i)$ as in definition 6.13, certifying that a shortest solution of Σ cannot induce $\mathcal{L}_i$.

Consider Λ a μ -minimal surface component. By definition, this means that the pruning process defined in section 7.6 stops. Let $(\Sigma', \mathcal{F}', \mu') = (\Sigma_{j_0}, \mathcal{F}_{j_0}, \mu_{j_0})$ the foliated complex obtained at the end of the pruning process. The Rips band complex $\Lambda'_{\mu'}$ is a (non-orientable) interval exchange in the following sense: every $x \in D_{\Lambda'_{\mu'}}$ outside a finite set belongs to exactly two bases of $\Lambda'_{\mu'}$ ([GLP94, Cor. 6.3]). Opening-up a separatrix (or unzipping a train-track carrying the foliation) gives a sequence of interval exchanges with arbitrary small domains. We want to implement this *unzipping process* at the level of Σ .

Consider $x_\mu \in D_{\Lambda'_{\mu'}}$ a point lying in 3 bands of $\Lambda'_{\mu'}$. Let $B_1 B_2 \dots B_j \dots$ be the infinite $\Lambda'_{\mu'}$ -word corresponding to the unique infinite regular leaf path starting at x_μ , and consider the corresponding Λ' -word $w_j = B_1 \dots B_j$.

Lemma 7.35. *There exists $x \in \pi_{\mu'}^{-1}(\{x_\mu\})$ lying in a singular leaf of $(\Sigma', \mathcal{F}')$, such that for all $j \geq 0$, $w_j(x)$ is not in the interior of a base of a band of $\Sigma' \setminus \Lambda'$.*

Proof. Let $I_1, \dots, I_p$ be the bases of the bands of $\Sigma' \setminus \Lambda'$ which lie in $D_{\Lambda'}$. Recall that $\mu'(I_1 \cup \dots \cup I_p) = 0$. Let

$$E = \{x \in \pi_{\mu'}^{-1}(\{x_\mu\}) \mid \exists k \text{ s.t. } h_{w_j}(x) \in I_1 \cup \dots \cup I_p\}.$$

Since the points $h_{w_j}(x_\mu)$ are all distinct in $\Lambda'_{\mu'}$, for each i and each $x \in \pi_{\mu'}^{-1}(x_\mu)$, there is at most one index k such that $w_j(x) \in I_i$. It follows that E is a finite union of segments. If E is empty, just take any point $x \in \pi_{\mu'}^{-1}(x_\mu)$ lying in a singular leaf of Σ . Otherwise, the leftmost point of E satisfies the Lemma. $\square$

We are now ready to describe the unzipping process. Consider $x \in D_{\Lambda'}$ as given by the lemma above and $x_j = h_{w_j}(x) \in D_{\Lambda'}$. For each $j \geq 1$, x_j lies in the interior of exactly two bases of Λ' . For $j = 0, 1, \dots$ perform the following moves on Σ' , modifying only Λ' :

- *Step a_j .* Cut the domain at x_j .
- *Step b_j .* Perform all possible forgetful moves on Λ' .

We denote by $(\Sigma_{i_0+j}, \mathcal{F}_{i_0+j}, \mu_{i_0+j})$ the band complex obtained after performing steps a_j and b_j . Proposition 7.34 will follow from the two following facts.

Lemma 7.36. *The complexity of Σ_j , defined as the sum of number of bands of Σ_j and of the number of connected components of D_j remains bounded.*

In particular, only finitely many distinct unfoliated band complexes with rational constraints appear in the sequence $\Sigma_1, \Sigma_2, \dots$.

Proof. Before step a_0 , x lies in the interior of at least 1 and at most 3 bases of Λ' because x_μ lies in the boundary of two bases and in the interior of one base of Λ'_μ . By choice of x , x does not lie in the interior of a base of band of $\Sigma' \setminus \Lambda'$. Step a_0 may increase $b_1(\Sigma)$ by at most 2. After step a_0 , x_1 lies in exactly 3 bases of Λ , and the interior of exactly one base by Lemma 7.35. This does not change after set b_0 .

By induction, we see that that steps a_j, b_j do not change $b_1(\Lambda)$, and after these steps, x_j lies in exactly 3 bases of Λ , and the interior of exactly one base by Lemma 7.35. As in the proof of Lemma 7.32, consider the graph Γ_j whose vertex set $V(\Gamma_j)$ is the set of connected components of D_{Λ_j} and whose set of non-oriented edges $E(\Gamma_j)$ corresponds to non-oriented bands. Since Λ_{j,μ_j} is an interval exchange, G_j has no vertex of valence one. Because of step b, each vertex of valence 2 of Γ_j is bounded by the number of bases of bands of $\Sigma_j \setminus \Lambda_j$. Since $b_1(\Gamma_j)$ is bounded, the lemma follows. $\square$

Lemma 7.37. *For all $\eta > 0$, there exists j large enough such that $\mu(D_{\Lambda_j}) < \eta$.*

Proof. Since $\{w_j(x_\mu)\}_{j \geq 0}$ is dense in $D_{\Lambda'_\mu}$, the maximum of the measures of the connected components of $D_{\Lambda'} \setminus \{x_0, \dots, x_j\}$ goes to 0 as j tends to infinity. It follows that the measure of the connected components of D_{Λ_j} go to 0 as j goes to infinity. Since the number of connected components of D_{Λ_j} is bounded, the lemma follows. $\square$

The proof of Proposition 7.34 is now identical to the proof of Proposition 7.31.

8 Endgame

8.1 Finding solutions of band complexes

Theorem 8.1. *There is an algorithm which takes as input a band complex with rational constraints, and decides whether it has a solution or not.*

Let us recall that we have defined two algorithms. The first one is the *prelamination generator*, described in Section 6.1. It takes as input a band complex Σ with rational constraints, and explores the space of all prelaminations (with rational constraints). The prelaminations are organised into a rooted tree $\mathcal{T}$ of finite valence, in such a way that children of a prelamination $\mathcal{L}$ are extensions of $\mathcal{L}$. If σ is a solution of Σ , it induces a Möbius-complete prelamination $\mathcal{L}_\infty(\sigma)$ (see Sections 5.1 and 5.2 for definitions). Lemma 6.2 says that $\mathcal{L}_\infty(\sigma)$ will be produced by the prelamination generator.

The second algorithm is the *prelamination analyser*, described in Subsection 6.2. It takes as input the band complex Σ together with a prelamination $\mathcal{L}$ (with rational constraints). If $\mathcal{L}$ is Möbius-complete, it decides if $\mathcal{L}$ is induced by a solution or not (see Lemma 6.4), and in this case, the prelamination analyser stops and outputs “Solution found” together with a solution, or “Reject” accordingly. If $\mathcal{L}$ is not Möbius-complete, it looks for some certificate ensuring that no shortest solution can induce

$\mathcal{L}$. If it can find one, it says “Reject” and stops. Otherwise, it may fail to stop or say “I don’t know”. The certificate is either

- the detection of an incompatibility of rational constraints,
- the detection of the non-existence of invariant measure,
- the detection of an exponent of periodicity that is too large for a shortest solution in regards of Bulitko’s Lemma (Prop. 4.1),
- or the detection of a shortening sequence of moves on $(\Sigma, \mathcal{L})$, which proves that a solution inducing $\mathcal{L}$ cannot be shortest.

See Section 6.2 for more details.

We can now explain the structure of the main algorithm.

Algorithm 8.2. Main algorithm

Input: a band complex with rational constraints Σ

Output: a solution of Σ , or “*There is no solution*”

Run the prelamination generator to construct step by step the rooted tree $\mathcal{T}$ of prelaminations

For each prelamination $\mathcal{L}$ produced by the generator, do in the background:

if $\mathcal{L}$ is a leaf of $\mathcal{T}$ and is not Möbius-complete **then**

 └ mark $\mathcal{L}$ as “*Rejected*”

 Run the prelamination analyser on $\mathcal{L}$

if the analyser says “Solution found” **then**

 └ **exit** with the solution provided by the analyser

if the analyser says “Reject” **then**

 └ mark $\mathcal{L}$ as “*Rejected*”

Simultaneously:

if some prelamination $\mathcal{L}$ is not a leaf and all children $\mathcal{L}$ get marked as “*Rejected*” **then**

 └ mark $\mathcal{L}$ as “*Rejected*”

if the root lamination get marked as “*Rejected*” **then**

 └ **exit** saying “*There is no solution*”

Remark. If some prelamination $\mathcal{L}$ is rejected by the prelamination analyser, we could also reject all its descendants, and we could ask the prelamination generator to stop extending this prelamination.

The algorithm could easily provide a proof that there is no solution when this is the case.

The main difficulty is to prove that the algorithm always stops. Theorem 8.1 follows immediately from the two following lemmas.

Lemma 8.3. *The main algorithm is correct: if it stops, its answer is true.*

Proof. By Lemma 6.15, the analyser is correct. Thus, the main algorithm is correct if it stops because the analyser has found a solution.

We claim that any prelamination which is marked as rejected cannot be induced by a shortest solution of Σ .

We prove the claim inductively. If $\mathcal{L}$ is marked as rejected by the prelamination analyser, then the claim is true by correctness of the prelamination analyser. If $\mathcal{L}$ is marked as rejected because $\mathcal{L}$ is a leaf of $\mathcal{T}$ and is not Möbius-complete, then

the claim is true because of Lemma 6.1. If $\mathcal{L}$ is not a leaf of $\mathcal{T}$ and if all its sons are correctly marked as rejected, then the claim holds for $\mathcal{L}$ by Corollary 6.2. This proves the claim.

If the root prelamination has been marked as rejected, then the root prelamination cannot be induced by a shortest solution of Σ , so Σ has no shortest solution, and has no solution at all. $\square$

Lemma 8.4. *The main algorithm always stops.*

Proof. Let Σ_0 be the band complex taken as input. If Σ_0 has a solution, then it has a shortest solution σ , and the prelamination $\mathcal{L}_\infty(\sigma)$ will be produced by the prelamination generator. If the algorithm does not stop, at some point $\mathcal{L}_\infty(\sigma)$ will be analysed by the analyser. Since $\mathcal{L}_\infty(\sigma)$ is Möbius-complete, the prelamination analyser will find out that there is a solution, and the main algorithm will stop, a contradiction.

Assume that Σ_0 has no solution. We claim that if some prelamination $\mathcal{L}$ is never rejected, then it is not a leaf, and at least one of its children is never rejected. Indeed, every leaf of the rooted tree $\mathcal{T}$ of prelaminations will be marked as rejected: either because it is not Möbius-complete, or because the analyser will reject it. On the other hand, if all children of $\mathcal{L}$ are marked as rejected at some point, then $\mathcal{L}$ will itself be marked as rejected. This proves the claim.

Assume by contradiction that the algorithm does not stop. Then the root lamination never gets rejected. The claim says that there is an infinite ray in $\mathcal{T}$ of prelaminations which are never rejected. This ray is an infinite sequence of prelaminations with rational constraints extending each other: $\mathcal{L}_1 \prec \mathcal{L}_2 \prec \dots$ (Definition 5.8).

First, there exists a topological foliation $\mathcal{F}_0$ on Σ_0 representing the prelamations $\mathcal{L}_i$ (this is by Proposition 7.3, see Definition 7.1 for the notion of foliation representing a sequence of prelaminations). Moreover, since no $\mathcal{L}_i$ is rejected for large exponent of periodicity, there exists a measure μ_0 , without atom, invariant under the holonomy of $\mathcal{F}_0$. This is ensured by Proposition 7.5.

Now by Proposition 7.21, if i is large enough, one can uniformly perform a finite sequence of inert moves on $(\Sigma_0, \mathcal{L}_i)$, and get a sequence of prelaminations represented in a foliated band complex $(\Sigma_1, \mathcal{F}_1, \mu_1)$ which is decomposed into μ -minimal components.

By Proposition 7.23, the Rips band complex $\Sigma_{1\mu_1}$ corresponding to $(\Sigma_1, \mathcal{F}_1, \mu_1)$ cannot contain a homogeneous minimal component, since otherwise, $\mathcal{L}_i$ would be rejected by the prelamination analyser for i large enough because of large exponent of periodicity.

Proposition 7.27 then says that one can uniformly perform a finite sequence of inert moves on $(\Sigma_1, \mathcal{F}_1, \mu_1)$ to get some band complex $(\Sigma, \mathcal{F}, \mu)$ whose bands are μ -independent. There are two alternatives left by the classification of minimal components: either there is an exotic component in $(\Sigma, \mathcal{F}, \mu)$, or there is a surface component. In both cases, there exists a shortening sequence of moves for $\mathcal{L}_i$ for i large enough (see Propositions 7.31 and 7.34). This proves the existence of shortening certificates for $\mathcal{L}_i$ for i large enough, contradicting the fact that no $\mathcal{L}_i$ is rejected. $\square$

8.2 Reduction of main theorems to band complexes

Proposition 8.5. *There exists an algorithm that takes as input a basis of a free group F , a finite set Φ of automorphisms of F preserving $S \cup S^{-1}$, and a system of twisted equations with rational constraints in F (with twisting automorphisms in Φ) and that decides whether there is a solution or not.*

Proof. By Proposition 3.1, this reduces to deciding whether a band complex has a solution. Theorem 8.1 concludes. $\square$

Theorem 3. *The problem of equations with rational constraints is solvable in finite extensions of free groups.*

More precisely, there exists an algorithm which takes as input a presentation of a virtually free group G , and a system of equations and inequations with constants in G , together with a set of rational constraints, and which decides if there exists a solution or not.

Proof. By proposition 2.4, the problem of equations with rational constraints in a virtually free group V reduces to the problem of twisted equations with rational constraints in the sense of Definition 2.2 where the twisting morphisms permute a basis. Proposition 8.5 concludes. $\square$

Without assuming that twisting morphisms preserve a basis of S , we can prove:

Theorem 4. *There exists an algorithm that takes as input a basis of a free group F , a finite set Φ of automorphisms of F whose image in $\text{Out}(F)$ generate a finite subgroup, and a system of twisted equations with rational constraints in F (with twisting automorphisms in Φ) and that decides whether there is a solution or not.*

Proof. Let $Q \subset \text{Out}(F)$ be the finite group generated by the image of Φ , and let V be the preimage of Q in $\text{Aut} F$. Embed F into $\text{Aut} F$ via inner automorphisms $x \mapsto i_x$ where $i_x(g) = xgx^{-1}$. Note that for all $\varphi \in \Phi$ and all $x \in F$, $i_{\varphi(x)} = \varphi \circ i_x \circ \varphi^{-1}$. It follows each equation twisted by Φ on F corresponds to a non-twisted equation in V , together with the rational constraint saying that the variables should lie in F . Since rational subsets of F are rational subsets of V , the theorem follows from Theorem 3. $\square$

9 Equations in hyperbolic groups with torsion

In [RS95], Rips and Sela constructed canonical representatives for a torsion free hyperbolic group Γ , in order to *lift* solutions of a system of equations on Γ to a free group. Then using Makanin's algorithm, they deduced an algorithm deciding whether a given system of equations in Γ has a solution.

Delzant [Del95, Rem.III.1] remarked that for certain hyperbolic groups with torsion, such canonical representatives do not exist, whereas Reinfeldt [Rei] noticed that, if the abelian subgroups are finite or cyclic, they do exist. In fact, in the general case, *most* of the construction of [RS95] remains valid, up to the construction of canonical cylinders. Instead of defining canonical representatives as paths in the Cayley graph of Γ (thus living in a free group) as in [RS95], we need to interpret them as paths in the 1-skeleton $\mathcal{K}$ of the barycentric subdivision of a Rips complex of Γ . The action of Γ on $\mathcal{K}$ fails to be free in presence of torsion, and paths in $\mathcal{K}$

correspond to elements of the virtually free group V occurring as the fundamental group of the graph of groups $\mathcal{K}/\Gamma$ (see below for details). The natural generalisation of Rips and Sela's construction in presence of torsion then leads to a family of systems of equations in the virtually free group V .

In [Dah05], rational constraints were used to handle inequations in torsion free hyperbolic groups. More generally, we would like to solve equations with rational constraints in a hyperbolic group. However, this is impossible without restriction on the rational subsets involved (even if Γ is torsion free). For instance, any finitely generated subgroup is a rational subset, but the membership problem is unsolvable in some hyperbolic groups ([Rip82]). This is why we introduce a nice class of rational subsets: *quasi-isometrically embeddable rational subsets* (see Definition 9.2 below). Examples of such subsets include finite subsets, quasiconvex subgroups, and their complements.

A set of *quasi-isometrically embeddable rational constraints* on a system of equations is the additional requirement that each variable x should live in a quasi-isometrically embeddable rational subset $\mathcal{R}_x \subset \Gamma$. As we will see, this class of subsets is a Boolean algebra (Cor. 9.6), so inequations are a particular case of quasi-isometrically embeddable rational constraints.

Here is the main statement of this section (compare with [Dah05, §5, 5.3]).

Theorem 1. *There exists an algorithm which takes as input*

- *a presentation $\langle S|R \rangle$ of a hyperbolic group Γ (maybe with torsion),*
- *a finite system of equations, inequations, with constants in Γ and with quasi-isometrically embeddable rational constraints*

and which decides whether there exists a solution or not.

Each rational constraint $\mathcal{R}_x$ should be given to the algorithm under the form of a finite automaton accepting a quasi-isometrically embedded language $\tilde{\mathcal{R}}_x \subset S_{\pm}^*$ projecting to $\mathcal{R}_x \subset \Gamma$ (see Definition 9.2).

Let us emphasise that the algorithm is uniform over all hyperbolic groups. This is a usual application of the fact that the hyperbolicity constant of a hyperbolic group can be computed from a presentation (see [Bow91, Prop 8.6.1] or [Gro87, Pap96]). In all the following, we will make sure that the algorithms we use are explicit if the hyperbolicity constant is known (in particular that the constants we use are explicit, or computable in terms of the presentation and the hyperbolicity constant). In this way, there is no restriction in considering that the hyperbolic group is given once and for all.

In particular, we get (compare with [Dah05, Theorem 0.1]):

Corollary 9.1. *The existential theory with constants of a hyperbolic group is decidable.*

9.1 Quasi-isometrically embeddable rational subsets

Definition 9.2. *Let Γ be a hyperbolic group generated by a finite set S , and $\pi : S_{\pm}^* \rightarrow \Gamma$ the corresponding morphism.*

A regular language $\tilde{\mathcal{R}} \subset S_{\pm}^$ is quasi-isometrically embedded in Γ if there exists $\lambda \geq 1, \mu \geq 0$, such that for any word $w \in \tilde{\mathcal{R}}$, $|\pi(w)|_{\Gamma} \geq \frac{1}{\lambda}|w| - \mu$.*

A rational subset $\mathcal{R} \subset \Gamma$ is quasi-isometrically embeddable in Γ if there exists a quasi-isometrically embedded regular language $\tilde{\mathcal{R}} \subset S_{\pm}^$ such that $\pi(\tilde{\mathcal{R}}) = \mathcal{R}$.*

Here, $|w|$ is the length of w as a word on $S_\pm$, and $|\cdot|_\Gamma$ is any word metric on Γ . In particular, finite sets and quasiconvex subgroups are clearly quasi-isometrically embeddable rational subsets.

The property of being quasi-isometrically embeddable does not depend on the chosen word metric on Γ . Neither does it depend on the choice generating set S . Indeed, consider any other generating set S' and express each element of S as a word on $S'_\pm$. This defines a morphism $\rho : S_\pm^* \rightarrow (S'_\pm)^*$ satisfying $|\rho(w)| \leq L|w|$ where $L = \max_{s \in S_\pm} |\rho(s)|$. Consider the natural morphism $\pi' : (S'_\pm)^* \rightarrow \Gamma$. The regular language $\tilde{\mathcal{R}}' = \rho(\tilde{\mathcal{R}})$ is quasi-isometrically embedded since for each $w' = \rho(w)$, $\pi'(w') = \pi(w)$, and $|\pi'(w')|_\Gamma = |\pi(w)|_\Gamma \geq \frac{1}{\lambda}|w| - \mu \geq \frac{1}{\lambda L}|w'| - \mu$.

However, the fact of being quasi-isometrically embedded does depend on the choice of $\tilde{\mathcal{R}}$ representing $\mathcal{R}$. For instance, if $\mathcal{R}$ is the set of all words, and $\mathcal{R}'$ is the set of L -local geodesics they represent the same rational subset (namely Γ), but $\mathcal{R}$ is not quasi-isometrically embedded.

A quasi-isometrically embeddable rational subset is quasiconvex: there exists $C > 0$ such that for all $x, y \in \mathcal{R}$, any geodesic of Γ joining x to y is contained in the C -neighbourhood of $\mathcal{R}$. Indeed, by thinness of triangles, one can assume $x = 1$. Let A be an automaton accepting $\tilde{\mathcal{R}}$. By stability of quasigeodesics, any path represented by a word accepted by $\tilde{\mathcal{R}}$ lies within a bounded distance from a geodesic joining its endpoints. Finally, if a path is accepted by A then any of its points lies a bounded distance away from an accepted point (the bound depending only on the number of states of A).

Conversely, we don't know if a quasiconvex rational subset is always quasi-isometrically embeddable.

9.2 Full sublanguage of a set of quasigeodesics

The goal of this section is to prove that given $\mathcal{R} \subset \Gamma$ a quasi-isometrically embeddable rational subset, the set of local quasigeodesics representing elements of $\mathcal{R}$ is rational (Proposition 9.4). This should be thought as an analogue of Lemma 1.6 saying that if $\langle S \rangle$ is a free group and $\mathcal{R} \subset \langle S \rangle$ is a rational subset, then the language of freely reduced words in $S_\pm^*$ representing an element of $\mathcal{R}$ is a regular language. As for the free group, we will deduce that quasi-isometrically embeddable rational subsets form a Boolean algebra.

Let S be a finite generating system of Γ , $\pi : S_\pm^* \twoheadrightarrow \Gamma$ be the corresponding morphism, $\text{Cay } \Gamma$ be the Cayley graph, and $|\cdot|_S$ be the word metric.

Given $w \in S_\pm^*$, we denote by p_w be the corresponding path in $\text{Cay } \Gamma$, defined by $p_w(i) = \pi(s_1 \dots s_i)$. Given $\lambda \geq 1$, $\mu \geq 0$, let $\mathcal{Q}\mathcal{G}_{\lambda, \mu}(S_\pm^*) \subset S_\pm^*$ be the set of (λ, μ) -quasigeodesics, i. e. the set of words w such that $|p_w(i) - p_w(j)|_S \geq \frac{1}{\lambda}|i - j| - \mu$. Similarly, the set of ν -local quasigeodesics $\mathcal{L}\mathcal{Q}\mathcal{G}_{\nu, \lambda, \mu}(S_\pm^*) \subset S_\pm^*$ is the set of words w such that for all $|i - j| \leq \nu$, $|p_w(i) - p_w(j)|_S \geq \lambda|i - j| - \mu$. Clearly, $\mathcal{L}\mathcal{Q}\mathcal{G}_{\nu, \lambda, \mu}(S_\pm^*)$ is a regular language since this is the complement of the set of words containing as a subword one of the finitely many words of length at most ν which are not (λ, μ) -quasigeodesic.

We will use the following statement about stability of local quasigeodesics:

Lemma 9.3 ([CDP90, Théorème 1.2, 1.4]). *Fix δ a hyperbolicity constant.*

Given $\lambda \geq 1$ and $\mu \geq 0$, there exists computable numbers ν , $\lambda' \geq 1$, $\mu' \geq 0$ such that any ν -local, (λ, μ) -quasigeodesic is a (λ', μ') -quasigeodesic.

Given $\lambda \geq 1, \mu \geq 0$, there exists a computable number η such that any (λ, μ) -quasigeodesic is at Hausdorff distance at most η from any geodesic with the same endpoints.

We always assume that ν is large enough so that all ν -local (λ, μ) -quasigeodesics are global quasigeodesics as in the result above.

Proposition 9.4. *Consider a hyperbolic group Γ , $\pi : S_{\pm}^* \rightarrow \Gamma$, and $\mathcal{R}$ a quasi-isometrically embeddable regular language. Consider any $\lambda \geq 1, \mu \geq 0$, and any ν such that ν -local (λ, μ) -quasigeodesics are global quasigeodesics.*

Then $\tilde{\mathcal{R}} = \pi^{-1}(\mathcal{R}) \cap \mathcal{LQ}\mathcal{G}_{\nu, \lambda, \mu}(S_{\pm}^)$ is a regular language of $S_{\pm}^*$.*

Moreover, given an automaton accepting a quasi-isometrically embedded regular language representing $\mathcal{R}$, one can algorithmically compute an automaton A accepting $\tilde{\mathcal{R}}$.

We say that $\tilde{\mathcal{R}}$ is *full* in $\mathcal{LQ}\mathcal{G}_{\nu, \lambda, \mu}(S_{\pm}^*)$ since any word in $\mathcal{LQ}\mathcal{G}_{\nu, \lambda, \mu}(S_{\pm}^*)$ representing an element of $\mathcal{R}$ lies in $\tilde{\mathcal{R}}$.

Remark 9.5. Recall that the set of geodesic words in a hyperbolic group is itself regular, by finiteness of Cannon's so-called cone types [Can84]. Using this fact, one could modify the proposition by substituting the set of global geodesic words to $\mathcal{LQ}\mathcal{G}_{\nu, \lambda, \mu}(S_{\pm}^*)$. However, we need to use local quasigeodesics for solving equations and inequations in hyperbolic groups.

Corollary 9.6. *The class of all quasi-isometrically embeddable rational subsets of a hyperbolic group Γ is a Boolean algebra.*

Moreover, this can be algorithmically computed: given automata accepting quasi-isometrically embedded regular languages representing $\mathcal{R}_1, \mathcal{R}_2$, one can compute automata accepting quasi-isometrically embedded regular languages representing $\mathcal{R}_1 \cup \mathcal{R}_2$, $\mathcal{R}_1 \cap \mathcal{R}_2$ and $\Gamma \setminus \mathcal{R}_1$.

Proof of the corollary. Clearly, the union of two quasi-isometrically embeddable rational subsets is quasi-isometrically embeddable. Let $\mathcal{R} \subset \Gamma$ be a quasi-isometrically embeddable rational subset. By Lemma 1.5, we need only to prove that $\Gamma \setminus \mathcal{R}$ is quasi-isometrically embeddable. Let $\mathcal{L} = \mathcal{LQ}\mathcal{G}_{\nu, 1, 0}(S_{\pm}^*)$ be the set of ν -local geodesics (for some ν large enough so that local geodesics are quasigeodesics). By Proposition 9.4, $\tilde{\mathcal{R}} = \pi^{-1}(\mathcal{R}) \cap \mathcal{L}$ is a regular language of $S_{\pm}^*$. Since regular languages of $S_{\pm}^*$ form a Boolean algebra, $\tilde{\mathcal{R}}' = \mathcal{L} \setminus \tilde{\mathcal{R}}$ is a regular language, it is quasi-isometrically embedded, and since $\pi(\mathcal{L}) = \Gamma$, $\pi(\tilde{\mathcal{R}}') = \Gamma \setminus \mathcal{R}$. $\square$

Proof of Proposition 9.4. Let $\tilde{\mathcal{R}}_0 \subset \mathcal{Q}\mathcal{G}_{\lambda_0, \mu_0}(S_{\pm}^*)$ be a quasi-isometrically embedded regular language representing $\mathcal{R}$. Consider a finite monoid $\mathcal{M}$, a morphism $\rho : S_{\pm}^* \rightarrow \mathcal{M}$, and $\text{Accept} \subset \mathcal{M}$ such that $\tilde{\mathcal{R}}_0 = \rho^{-1}(\text{Accept})$. Let $\mathcal{P}$ be the language of prefixes of $\tilde{\mathcal{R}}_0$: $w \in \mathcal{P}$ if there exists $u \in S_{\pm}^*$ with $wu \in \tilde{\mathcal{R}}_0$. Note that $\mathcal{P}$ consists of (λ_0, μ_0) -quasigeodesics, and $\mathcal{P} = \rho^{-1}(P)$ where $P = \{m \in \mathcal{M} \mid \exists n \in \mathcal{M}, mn \in \text{Accept}\}$.

Write $\mathcal{L} = \mathcal{LQ}\mathcal{G}_{\nu, \lambda, \mu}(S_{\pm}^*)$. Let λ', μ' and η be such that ν -local (λ, μ) -quasigeodesics of Γ are global (λ', μ') -quasigeodesic, and is at Hausdorff distance at most η from any geodesic with the same endpoints (those values can be algorithmically computed from ν, λ, μ).

On the other hand, from the automaton A_0 , one can algorithmically find (λ'_0, μ'_0) such that all accepted words are (λ'_0, μ'_0) -quasigeodesics. Indeed, given any positive ν_1 , and any λ_1, μ_1 , one can algorithmically check whether all words accepted by

A_0 are ν_1 -local (λ_1, μ_1) -quasigeodesics. Denote by $\nu(\lambda, \mu)$, $\lambda'(\lambda, \mu)$ and $\mu'(\lambda, \mu)$ computable functions such that $\nu(\lambda, \mu)$ -local (λ, μ) -quasigeodesics are global $(\lambda'(\lambda, \mu), \mu'(\lambda, \mu))$ -quasigeodesics. By checking if $\tilde{\mathcal{R}}_0 \subset \mathcal{LQ}\mathcal{G}_{\nu(\lambda_1, \mu_1), \lambda_1, \mu_1}$ for larger and larger values of λ_1, μ_1 , one will finally get a positive answer, from which one can conclude that $\tilde{\mathcal{R}}_0 \subset \mathcal{Q}\mathcal{G}_{\lambda'(\lambda_1, \mu_1), \mu'(\lambda_1, \mu_1)}$. Compute η_0 such that each (λ'_0, μ'_0) -quasigeodesic is at Hausdorff distance at most η_0 from any geodesic with the same endpoints.

For $h \in \Gamma$, consider $\sigma_h = \rho(\mathcal{P} \cap \pi^{-1}(h)) \subset \mathcal{M}$ (σ_h can be thought as the set of states of the automaton corresponding to words representing h , and which can be extended to accepted words). Note that $h \in R$ if and only if $\sigma_h \cap \text{Accept} \neq \emptyset$.

Given $w \in \mathcal{L}$, Let p_w be the path in Cay Γ corresponding to w . We want to get a local picture of the values of σ_h in the neighbourhood of the endpoint $g = \pi(w)$ of p_w . Define $R = \eta_0 + \eta + 10\delta$ and $L = \max\{(R + 1 + 10\delta + \mu' + \eta)\lambda', \nu\}$. Consider w_L the suffix of length L of w ($w = w_L$ if $|w| \leq L$). Let q_w be the suffix of length L of the path $g^{-1}p_w$ (q_w is the path ending at 1 and labelled by w_L). Let N be the R -neighbourhood of q_w , and consider $\Sigma_w : N \rightarrow 2^{\mathcal{M}}$ defined by $\Sigma_w(h) = \sigma_{gh}$ (this records the values of σ_h in the neighbourhood of the suffix of p_w). In particular, $\pi(w) \in \mathcal{R}$ if and only if $\Sigma_w(1) \cap \text{Accept} \neq \emptyset$. The mapping $\Phi : w \mapsto (w_L, \Sigma_w)$ clearly takes finitely many values.

Claim. For all word $w \in S_{\pm}^*$ and all $s \in S_{\pm}$ with $w, ws \in \mathcal{L}$, $\Phi(ws)$ depends only on $\Phi(w)$ and s .

Proof. If $|w_L| < L$, this is clear as $\Phi(w)$ then determines $w = w_L$. Obviously, the L -suffix of ws is determined by w_L and s . We need to prove that for all $|h| \leq R$, Σ_w and w_L determine σ_{gsh} . Write w as a concatenation $w = w'.w_L$ and let $x_L = \pi(w') \in \Gamma$. Consider $u \in \mathcal{P}$ be a word with $\pi(u) = gsh$, and let p_u be the corresponding path in Cay Γ .

Fact. p_u intersects $B(x_L, R)$.

Proof of the fact. Consider c_u (resp. c_w) a geodesic joining 1 to gsh (resp. to g). The Hausdorff distance between c_u and p_u (resp. between c_w and p_w) is at most η_0 (resp. η). The projection x_w of x_L on c_w satisfies $d(x_w, x_L) \leq \eta$. Thus, $d(g, x_w) \geq -\eta + \frac{1}{\lambda'}L - \mu' \geq R + 1 + 10\delta$ by choice of L . Looking at a comparison tree for the triangle $1, g, gsh$, since $d(g, gsh) \leq R + 1$, we see that x_w is 10δ -close to some point $x_u \in c_u$. The projection x'_u of x_u on p_u is η_0 close to x_u and satisfies $d(x'_u, x_L) \leq \eta_0 + 10\delta + \eta = R$. $\square$

We now prove that the fact implies our claim. Given $u \in \mathcal{P}$ with $\pi(u) = gsh$, write $u = u'u''$ with $\pi(u') = x' \in B(x_L, R)$, and let $m' = \rho(u')$. Consider $y_L = g^{-1}x_L$ the initial point of q_w , and $y' = g^{-1}x' \in N$. For each $x' \in B(x_L, R)$, the set of possible m' is known as it is encoded in $\sigma_{x'} = \sigma_{gy'}$. Since u'' is a quasigeodesic between two points at distance at most $L + 2R + 1$, $|u''|$ is bounded by $L_{u''} = \lambda(L + 2R + 1 + \mu)$. Since $u \in \mathcal{P}$, among those quasigeodesic u'' , one should consider only those such that $m'\rho(u'') \in P$ i. e. $u'' \in \rho^{-1}(m'^{-1}P)$. Then σ_{gsh} is precisely the set of all possible values of $m'\rho(u'')$ as y' varies in $B(y_L, R)$, m' varies in $\sigma_{gy'}$ and u'' varies in the set of words of length at most $L_{u''}$ in $\rho^{-1}(m'^{-1}P)$. This proves the claim, and proves moreover that $\Phi(ws)$ is algorithmically computable from $\Phi(w)$ and s . $\square$

We now construct a deterministic automaton A accepting $\mathcal{L} \cap \pi^{-1}(\mathcal{R})$. Note that for any individual $w \in S_{\pm}^*$, $\Phi(w)$ is algorithmically computable since one can enumerate the finitely many (λ'_0, μ'_0) -quasigeodesics with endpoint $\pi(w)$.

Let $\mathcal{L}_{\leq L}$ be the set of words of length at most L in $\mathcal{L}$. Since the R -neighbourhood N of q_w satisfies $N \subset B(1, R+L)$, Σ_w is a partially defined map on $B(1, R+L)$, i. e. an element of the finite set $D = (2^{\mathcal{M}} \cup \{\text{undef}\})^{B(1, R+L)}$. Thus, $\Phi(\mathcal{L})$ takes values in the finite set $F = \mathcal{L}_L \times D$.

We take F as the set of states of our automaton, and $\Phi(1)$ as its initial state. If $f = (w_L, \Sigma_w) \in F$ with $|w_L| < L$, we can discard f if $f \neq \Phi(w_L)$. If $|w_L| < L$ and $f = \Phi(w_L)$, for each $s \in S_{\pm}$, we add an edge joining f to $\Phi(w_L s)$. If $|w_L| = L$, and if $w_L.s \in \mathcal{L}$, we connect (w_L, Σ_w) to the state (w', Σ') determined by (w_L, Σ_w) and s as in the claim. The set of words w read by the automaton starting from the initial state is exactly $\mathcal{L}$. Moreover, the corresponding final state is $\Phi(w)$ which determines $\sigma_{\pi(w)} = \Sigma_w(1)$. We define the set of accepting states of A as the set of elements $(w_L, \Sigma_w) \in F$ such that $\sigma_{\pi(w)} \cap \text{Accept} \neq \emptyset$. Since $\pi(w) \in \mathcal{R}$ if and only if $\Sigma_{\pi(s)} \cap \text{Accept} \neq \emptyset$, the language accepted by A is $\pi^{-1}(\mathcal{R}) \cap \mathcal{L}$.

Finally, note that the automaton A can be algorithmically computed from A_0 . $\square$

9.3 Canonical representatives in hyperbolic groups with torsion

9.3.1 Canonical sliced cylinders and paths.

Let Γ be a hyperbolic group, and $\text{Cay } \Gamma$ be a Cayley graph. Let δ be its hyperbolicity constant. A *cylinder* for $(x, y) \in \Gamma \times \Gamma$ is a subset of the 5δ -neighbourhood of a geodesic segment $[x, y]$ in $\text{Cay } \Gamma$, containing every such segment. A *slicing* of a cylinder (or a slice decomposition), is a partition into subsets (called slices), with a total ordering of them, such that

- the union of any two consecutive slices has diameter at most 50δ
- for this ordering, the slice containing x is smaller or equal to the slice containing y
- slices move quasigeodesically: if x, y lie respectively in the i -th and j -th slice for this ordering, then $d(x, y) \geq \frac{1}{4m_0}|i - j| - 200\delta$ where $m_0 = \#B(1, 50\delta)$.

In [RS95], Rips and Sela proved the following theorem (which applies for hyperbolic groups with torsion).

Theorem 9.7 ([RS95, Cor. 4.3]). *Let Γ be a hyperbolic group and $\text{Cay } \Gamma$ a Cayley graph. Then there exists a computable constant κ (controlling the size of the defect C_i below) such that the following holds.*

Let X be a set of variables, and $\mathcal{E}$ be a triangular system of equations over X . Let $\underline{g} = (g_x)_{x \in X \cup \overline{X}} \in \Gamma^X$ be a solution of $\mathcal{E}$.

Then for every $x \in X \cup \overline{X}$, there exists a cylinder Cyl_x for $(1, g_x)$ together with a slicing, such that

- *for each variable $x \in X$, $\text{Cyl}_{\overline{x}} = g_x^{-1}\text{Cyl}_x$, their slices are the same but in reverse order,*

- for each equation $x_1x_2x_3 \in \mathcal{E}$, and for each $i \in \{1, 2, 3 \bmod 3\}$ there is a decomposition (depending on the equation) of the cylinder as

$$\text{Cyl}_{x_i} = L_i \sqcup C_i \sqcup R_i$$

where

- L_i, C_i, R_i are union of slices with $L_i < C_i < R_i$ with respect to the ordering,
- $L_{i+1} = g_{x_i}^{-1} R_i$, their slices are the same but in reverse order,
- C_i contains at most $\kappa \cdot \#\mathcal{E}$ slices.




Figure 6: Three cylinders for a solution to $x_1x_2x_3 = 1$. The subdivision in slices is drawn. The central part, where the cylinders do not coincide, does not contain more than $3\kappa\#\mathcal{E}$ slices.

Note that cylinders and slicings depend only on the variable considered whereas the decomposition of the cylinder depends on the equation in which the variable appears. The fact that slices move quasigeodesically is a consequence of their definition in [RS95], see [Dah05, Prop. 3.8] for a proof.

Next step consists in interpreting sliced cylinders as paths in a suitable graph $\mathcal{K}$. Let us consider the Rips complex $P_{50\delta}(\Gamma)$ whose set of vertices is Γ , and whose simplices are subsets of Γ of diameter at most 50δ . Let us call $\mathcal{K}$ the 1-skeleton of its barycentric subdivision. We identify Γ to a subset of the vertices of $\mathcal{K}$, in the obvious way. For any path $\mathbf{p} : \{0, \dots, n\} \rightarrow \mathcal{K}$, we denote by $\overline{\mathbf{p}}$ the reverse path defined by $\overline{\mathbf{p}}(i) = \mathbf{p}(n - i)$. If $\mathbf{q}$ is a path with $\mathbf{q}(0) = \mathbf{p}(n)$, we denote by $\mathbf{p} \cdot \mathbf{q}$ the concatenation.

Proposition 9.8. *Let Γ be a hyperbolic group, $\text{Cay } \Gamma$ a Cayley graph, and $\mathcal{K}$ the 1-skeleton of the barycentric subdivision of the Rips complex of Γ as above. Consider $\lambda_0 = 400\delta m_0$ where m_0 is a bound on the cardinality of balls of radius 50δ in $\text{Cay } \Gamma$, $\mu_0 = 8$, and κ as in Theorem 9.7.*

Let $\mathcal{E}$ be a triangular system of equations over X , and $\underline{g} = (g_x)_{x \in X} \in \Gamma^X$ be a solution of $\mathcal{E}$ and let $g_{\overline{x}} = g_x^{-1}$.

Then for every $x \in X \cup \overline{X}$, there exists a (λ_0, μ_0) -quasigeodesic path $\mathbf{p}_x$ joining 1 to g_x in $\mathcal{K}$ such that

- for each variable $x \in X$, $\mathbf{p}_x = g_x^{-1} \overline{\mathbf{p}}_x$
- for each equation $x_1 x_2 x_3 \in \mathcal{E}$, and for each $i \in \{1, 2, 3 \bmod 3\}$ there is a decomposition (depending on the equation) $\mathbf{p}_{x_i} = \mathbf{l}_i \cdot \mathbf{c}_i \cdot \mathbf{r}_i$, where $\mathbf{l}_{i+1} = g_{x_i}^{-1} \overline{\mathbf{r}}_i$, and $\mathbf{c}_i$ has length at most $2\kappa \cdot \#\mathcal{E}$.

Proof. Consider sliced cylinders C_x given by Rips and Sela's Theorem 9.7. Let S, S' be the slices containing 1 and g_x respectively, and $S = S_1, \dots, S_n = S'$ be the set of slices between S and S' , in increasing order. Since S_i of C_x has diameter at most 50δ , it defines a simplex of $P_{50\delta}(\Gamma)$, and thus a vertex v_i of $\mathcal{K}$. Similarly, $S_i \cup S_{i+1}$ defines a vertex u_i of $\mathcal{K}$ connected by an edge to v_i and v_{i+1} , and there is an edge in $\mathcal{K}$ joining 1 to v_1 and v_n to g_x . We can therefore define $\mathbf{p}_x$ as $1, v_1, u_1, v_2, \dots, u_{n-1}, v_n, g_x$.

Let us explain why these paths are quasigeodesic, the other properties immediately follow from the properties of canonical cylinders. The fact that slices vary quasigeodesically say that the distance in $\text{Cay } \Gamma$ between the i -th slice and the j -th slice is at least $|j - i|/4m_0 - 200\delta$. Thus, the barycentres of the corresponding simplices in $P_{50\delta}(\Gamma)$ are at distance at least $|j - i|/(200\delta m_0) - 4$ so $d_{\mathcal{K}}(v_i, v_j) \geq |j - i|/(200\delta m_0) - 4$. Taking into account the fact that $\mathbf{p}_x$ takes $2|j - i|$ steps to go from v_i to v_j , we get $d_{\mathcal{K}}(\mathbf{p}_x(s), \mathbf{p}_x(t)) \geq |s - t|/(400\delta m_0) - 4$ for all odd s, t . Since u_i is at distance 1 from v_i and v_{i+1} , one gets for all s, t $d_{\mathcal{K}}(\mathbf{p}_x(s), \mathbf{p}_x(t)) \geq (|s - t| - 2)/(400\delta m_0) - 4 - 2 \geq |s - t|/(400\delta m_0) - 8$ as desired. $\square$

9.3.2 The virtually free group V , and canonical representatives for Γ .

We now define the virtually free group V as a group of paths. We consider homotopy classes of edge paths relative to endpoints in the graph $\mathcal{K}$. A path is *reduced* if it has no subpath of length 2 passing twice through the same edge. Thus, each homotopy class of paths in $\mathcal{K}$ has a unique reduced representative.

Let V be the set of all homotopy classes of edge paths $\mathbf{p}$ in $\mathcal{K}$ that start at the vertex 1_Γ , and end at a vertex of Γ . Define $\pi : V \rightarrow \Gamma$ by mapping $\mathbf{p}$ to its endpoint. We endow V with a group structure by defining $\mathbf{p}\mathbf{p}'$ to be the homotopy class of the concatenation $\mathbf{p} \cdot (\pi(\mathbf{p})\mathbf{p}')$ (where $\pi(\mathbf{p})\mathbf{p}'$ is the translate of $\mathbf{p}'$ by $\pi(\mathbf{p}) \in \Gamma$). Note that $\mathbf{p}^{-1} = \pi(\mathbf{p})^{-1}\overline{\mathbf{p}}$. It is also clear that $\pi : V \rightarrow \Gamma$ is a surjective homomorphism.

Lemma 9.9. *The group V is virtually free. More precisely, it is isomorphic to the fundamental group of the finite graph of finite groups defined by $\mathcal{K}/\Gamma$, where vertices and edges are marked by copies of stabilisers of their preimages in $\mathcal{K}$. In particular, a presentation of V is computable from a presentation of Γ .*

Proof. Let $p : T \rightarrow \mathcal{K}$ be the universal cover of the graph $\mathcal{K}$; it is a tree. With the choice of a preimage $v_0 \in T$ of $1_\Gamma \in \mathcal{K}$, the group V admits a natural action on T , as we explain now. Any path $\mathbf{p}$ in $\mathcal{K}$ starting at 1_Γ has a unique lift $\tilde{\mathbf{p}}$ in T starting at v_0 . On the other hand, for any point $x \in T$, the path $[v_0, x] \subset T$ defines a path $\mathbf{q}_x$ by projection to $\mathcal{K}$. Then one can define $\mathbf{p}.x$ as the endpoint of lift starting at v_0 of $\mathbf{p} \cdot (\pi(\mathbf{p})\mathbf{q}_x)$. It is easy to check that $p : T \rightarrow \mathcal{K}$ is π -equivariant. In particular, $T/V \simeq \mathcal{K}/\Gamma$ is a finite graph.

Note that $\pi_1(\mathcal{K}, 1) \subset V$ is the kernel of π and acts freely on T .

To show that V is virtually free, we prove that stabilisers of vertices of T are finite. Let $V_x \subset V$ be the stabiliser of a vertex $x \in T$. Then $\pi(V_x)$ stabilises

$p(x) \in \mathcal{K}$. Since $\ker \pi$ acts freely on T , $\pi|_{V_x}$ is injective. This proves the finiteness of vertex stabilisers.

To prove the computability of vertex stabilisers, we prove that π induces an isomorphism between V_x and the Γ stabiliser of $p(x)$, which is computable. We need to prove that any $g \in \Gamma$ fixing $p(x)$ lies in $\pi(V_x)$. Consider the path $\mathbf{q}' = g\mathbf{q}_x$ joining g to $p(x)$, and $\mathbf{p} = \mathbf{q}_x \cdot \overline{\mathbf{q}'}$ joining 1 to g . Then $\mathbf{p} \cdot x$ is the endpoint of the lift of $\mathbf{p} \cdot (g\mathbf{q}_x) = \mathbf{q}_x \cdot \overline{\mathbf{q}'} \cdot (g\mathbf{q}_x) = \mathbf{q}_x$, so $\mathbf{p} \cdot x = x$.

A similar argument shows that π induces an isomorphism between edge stabilisers of T and $\mathcal{K}$, so one can compute a finite graph of finite groups representing V . $\square$

9.4 Lifting equations to V

Consider $(\lambda_1, \mu_1) = (\lambda_0, \mu_0 + 2 + \frac{2}{\lambda_0})$, so that any concatenation of a (λ_0, μ_0) -quasigeodesic with a path of length 1 at each extremity is a (λ_1, μ_1) -quasigeodesic. Let $\mathcal{QG}_{\lambda_1, \mu_1}(V) \subset V$ be the set of elements such that the corresponding reduced path in $\mathcal{K}$ is (λ_1, μ_1) -quasigeodesic. Denote by $V_{\leq L}$ the set of elements of V whose corresponding reduced path in $\mathcal{K}$ has length at most L .

Interpreting the paths occurring in Proposition 9.8 in terms of elements of V , we get:

Proposition 9.10. *Consider a system of equations in Γ as in Proposition 9.8, and $(g_x)_{x \in X} \in \Gamma^X$ a solution. Let $\kappa_1 = 2\kappa \cdot \#\mathcal{E} + 2$.*

Then for each variable $x \in X \cup \overline{X}$, there exists $\tilde{g}_x \in \mathcal{QG}_{\lambda_1, \mu_1}(V)$ with $\tilde{g}_{\overline{x}} = (\tilde{g}_x)^{-1}$ and $\pi(\tilde{g}_x) = g_x$, and for each equation $\varepsilon \in \mathcal{E}$ representing the equation $x_1 x_2 x_3 = 1$ and for each $i \in \{1, 2, 3 \bmod 3\}$, there exists $\tilde{l}_{\varepsilon, i} \in \mathcal{QG}_{\lambda_1, \mu_1}(V)$ and $\tilde{c}_{\varepsilon, i} \in V_{\leq \kappa_1}$, such that

- (i) $\tilde{g}_{x_i} = \tilde{l}_{\varepsilon, i} \tilde{c}_{\varepsilon, i} \tilde{l}_{\varepsilon, i+1}^{-1}$ in V
- (ii) $\pi(\tilde{c}_{\varepsilon, 1} \tilde{c}_{\varepsilon, 2} \tilde{c}_{\varepsilon, 3}) = 1$ in Γ .

Conversely, given any family of elements of V $(\tilde{g}_x)_{x \in X \cup \overline{X}}$, $(\tilde{l}_{\varepsilon, i})_{\varepsilon \in \mathcal{E}, i=1,2,3}$, $(\tilde{c}_{\varepsilon, i})_{\varepsilon \in \mathcal{E}, i=1,2,3}$ satisfying $\tilde{g}_{\overline{x}} = \tilde{g}_x^{-1}$ and (i) – (ii), the family $g_x = \pi(\tilde{g}_x)$ is a solution of $\mathcal{E}$.




Figure 7: canonical paths in $\mathcal{K}$

Proof. The converse implication is obvious as for each equation ε written $x_1 x_2 x_3 = 1$, $\pi(\tilde{g}_{x_1} \tilde{g}_{x_2} \tilde{g}_{x_3})$ simplifies to $\pi(\tilde{l}_{\varepsilon,1} \tilde{c}_{\varepsilon,1} \tilde{c}_{\varepsilon,2} \tilde{c}_{\varepsilon,3} \tilde{l}_{\varepsilon,1}^{-1}) = 1$.

For the direct implication, apply Proposition 9.8 to get a path $\mathbf{p}_x$ for each $x \in X \cup \overline{X}$, and for each equation ε written $x_1 x_2 x_3 = 1$, and each $i \in \{1, 2, 3 \bmod 3\}$, three paths $\mathbf{l}_{\varepsilon,i}, \mathbf{c}_{\varepsilon,i}, \mathbf{r}_{\varepsilon,i}$. Since $\mathbf{p}_x$ joins 1 to g_x , it represents an element $\tilde{g}_x = \mathbf{p}_x \in V$, and $\pi(\tilde{g}_x) = g_x$. On the other hand $\mathbf{l}_{\varepsilon,i}$ may fail to end at a vertex of Γ , so consider $e_{\varepsilon,i}$ an edge of $\mathcal{K}$ joining the endpoint of $\mathbf{l}_{\varepsilon,i}$ to a point in $l_{\varepsilon,i}\Gamma$. This allows to define an element of V by $\tilde{l}_{\varepsilon,i} = \mathbf{l}_{\varepsilon,i} \cdot e_{\varepsilon,i}$. Since $\mathbf{l}_{\varepsilon,i+1} = g_{x_i}^{-1} \overline{\mathbf{r}_{\varepsilon,i}}$, the edge $g_{x_i} e_{\varepsilon,i+1}$ joins the initial point of $\mathbf{r}_{\varepsilon,i}$ to $g_{x_i} l_{\varepsilon,i+1} \in \Gamma$, we denote it by $e'_{\varepsilon,i}$ (see Figure 7). One can therefore define the following elements of V : $\tilde{c}_{\varepsilon,i} = l_{\varepsilon,i}^{-1}(\overline{e_{\varepsilon,i}} \cdot \mathbf{c}_{\varepsilon,i} \cdot e'_{\varepsilon,i})$, and $\tilde{r}_{\varepsilon,i} = (g_{x_i} l_{\varepsilon,i+1})^{-1}(\overline{e'_{\varepsilon,i}} \cdot \mathbf{r}_{\varepsilon,i})$.

By construction, $\tilde{g}_{x_i} = \tilde{l}_{\varepsilon,i} \tilde{c}_{\varepsilon,i} \tilde{r}_{\varepsilon,i}$. Since $\mathbf{l}_{\varepsilon,i+1} = g_{x_i}^{-1} \overline{\mathbf{r}_{\varepsilon,i}}$, we get $\tilde{l}_{\varepsilon,i+1} = \mathbf{l}_{\varepsilon,i+1} \cdot e_{i+1} = g_{x_i}^{-1}(\overline{\mathbf{r}_{\varepsilon,i}} \cdot e'_{\varepsilon,i}) = \tilde{r}_{\varepsilon,i}^{-1}$. The fact that $\pi(\tilde{c}_{\varepsilon,1} \tilde{c}_{\varepsilon,2} \tilde{c}_{\varepsilon,3}) = 1$ then follows from $\pi(\tilde{g}_{x_1} \tilde{g}_{x_2} \tilde{g}_{x_3}) = 1_\Gamma$. The fact that $\tilde{l} \in \mathcal{QG}_{\lambda_1, \mu_1}(V)$ and $\tilde{c} \in V_{\leq \kappa_1}$ is clear from the analogous facts satisfied by $\mathbf{l}$ and $\mathbf{c}$. $\square$

Note that statement (i) of Proposition 9.10 is an equation in V while (ii) takes place in Γ . But since each $\tilde{c}_{\varepsilon,i}$ is short, there are only finitely possibilities, and for each value of $\tilde{c}_{\varepsilon,i}$ satisfying (ii), one can think of (i) as a system equations in V with variables $\tilde{l}_{\varepsilon,i}$ and $\tilde{g}_x$, and constants $\tilde{c}_{\varepsilon,i}$. More formally:

Definition 9.11. For each tuple $\tilde{c}$ of elements $\tilde{c}_{\varepsilon,i} \in V_{\leq \kappa_1}$ satisfying (ii), define $\tilde{\mathcal{E}}(\tilde{c})$ as the system of equations (i) with unknowns $\tilde{g}_x$ and $\tilde{l}_{\varepsilon,i}$.

For each solution of $\tilde{\mathcal{E}}(\tilde{c})$, its projection $(g_x) = (\pi(\tilde{g}_x)) \in \Gamma^X$, is a solution of $\mathcal{E}$. We say that the solution $(\tilde{g}_x, \tilde{l}_{\varepsilon,i})$ is a lift of (g_x) in V .

Proposition 9.10 says that every solution of $\mathcal{E}$ has a lift in V , and more precisely in $\mathcal{QG}_{\lambda_1, \mu_1}(V) \subset V$.

9.5 Lifting rational constraints to V , and proof of the theorem

Before proving the main theorem of this section, we need to lift the quasi-isometrically embedded rational subsets of Γ in V . Let $\hat{S}$ be a generating system of V . Let $\mathcal{L}_{\hat{S}} = \mathcal{LQG}_{\nu, \lambda, \mu} \subset \hat{S}_\pm^*$ be the the rational language of local quasigeodesics for the metric $|\cdot|_{\hat{S}}$ on Γ . Let $\mathcal{L}_V$ be its image in V . Given λ_1, μ_1 as in previous section, one can compute λ, μ large enough so that $\mathcal{L}_V$ contains $\mathcal{QG}_{\lambda_1, \mu_1}(V)$.

Lemma 9.12. Let $\mathcal{R}$ be any quasi-isometrically embeddable rational subset of a hyperbolic group G . Consider V , $\pi : V \twoheadrightarrow G$ and $\mathcal{QG}_{\lambda_1, \mu_1}(V) \subset \mathcal{L}_V \subset V$ as above.

Then $\tilde{\mathcal{R}}_V = \mathcal{L}_V \cap \pi^{-1}(\mathcal{R})$ is a rational subset of V , and $\pi(\tilde{\mathcal{R}}_V) = \mathcal{R}$.

Proof. By Proposition 9.4, the language $\tilde{\mathcal{R}}_{\hat{S}} = \pi^{-1}(\mathcal{R}) \cap \mathcal{L}_{\hat{S}} \subset \hat{S}_\pm^*$ is regular. Since $\pi_{\hat{S}} : \hat{S}_\pm^* \rightarrow G$ factors through $\pi_V : V \rightarrow G$, $\tilde{\mathcal{R}}_V$ is the image in V of $\tilde{\mathcal{R}}_{\hat{S}}$. Therefore, it is rational. $\square$

We are now ready to prove that one can decide algorithmically whether a system of equations with quasi-isometrically -embeddable rational constraints has a solution.

Proof of Theorem 1. Since $\Gamma \setminus \{1\}$ is a quasi-isometrically embeddable rational subset by Corollary 9.6, and since constants can be encoded by rational constraints consisting of a single element, we can reduce to the case of a system of equations

$\mathcal{E}$ with quasi-isometrically embeddable rational constraints $\mathcal{R}_x \subset \Gamma$ (without constants).

For each $x \in X$, consider $\tilde{\mathcal{R}}_x = \mathcal{L}_V \cap \pi^{-1}(\mathcal{R}_x) \subset V$ which is rational by Lemma 9.12. Compute κ_1 , and enumerate all tuples $\tilde{c}$ in $V_{\leq \kappa_1}$ satisfying $\pi(\tilde{c}_{\varepsilon,1}\tilde{c}_{\varepsilon,2}\tilde{c}_{\varepsilon,3}) = 1$ (this uses a solution of the word problem in Γ). Consider the lifted system of equations $\tilde{\mathcal{E}}(\tilde{c})$ in V as in definition 9.11, and add the rational constraint $\tilde{g}_x \in \tilde{\mathcal{R}}_x$. Denote by $\tilde{\mathcal{R}}$ the corresponding set of rational constraints. For each $\tilde{c}$, use the algorithm of Theorem 3 for virtually free groups to decide whether $(\tilde{\mathcal{E}}(\tilde{c}), \tilde{\mathcal{R}})$ has a solution. We claim that $(\mathcal{E}, \mathcal{R})$ has a solution if and only if $(\tilde{\mathcal{E}}(\tilde{c}), \tilde{\mathcal{R}})$ has a solution for at least one $\tilde{c}$.

If $(\tilde{\mathcal{E}}(\tilde{c}), \tilde{\mathcal{R}})$ has a solution, then $\pi(\tilde{g}_x)$ is a solution of $\mathcal{E}$ by Proposition 9.10, and since $\tilde{g}_x \in \tilde{\mathcal{R}}_x$, $\pi(\tilde{g}_x) \in \mathcal{R}_x$, so the constraints are satisfied.

Conversely, if $(\mathcal{E}, \mathcal{R})$ has a solution (g_x) , then by Proposition 9.10, $\tilde{\mathcal{E}}(\tilde{c})$ has a solution lying in $\mathcal{QG}_{\lambda_1, \mu_1}(V) \subset \mathcal{L}_V$ for some tuple $\tilde{c}$. Since the solution satisfies the rational constraint $g_x \in \mathcal{R}_x$, the corresponding solution of $\tilde{\mathcal{E}}(\tilde{c})$ satisfies $\tilde{g}_x \in \mathcal{L}_V \cap \pi^{-1}(\mathcal{R}_x) = \tilde{\mathcal{R}}_x$, so this solution of $\tilde{\mathcal{E}}(\tilde{c})$ satisfies the rational constraint $\tilde{\mathcal{R}}$. $\square$

References

- [Ber79] Jean Berstel. *Transductions and context-free languages*, volume 38 of *Leitfäden der Angewandten Mathematik und Mechanik [Guides to Applied Mathematics and Mechanics]*. B. G. Teubner, Stuttgart, 1979.
- [BF95] Mladen Bestvina and Mark Feighn. Stable actions of groups on real trees. *Invent. Math.*, 121(2):287–321, 1995.
- [BMS] Gilbert Baumslag, Alexei Myasnikov, and Vladimir Shpilrain. Open problems in combinatorial group theory. <http://web.cs.gc.cuny.edu/~cryptlab/gworld/problems/oproblems.html>.
- [Bow91] B. H. Bowditch. Notes on Gromov’s hyperbolicity criterion for path-metric spaces. In *Group theory from a geometrical viewpoint (Trieste, 1990)*, pages 64–167. World Sci. Publishing, River Edge, NJ, River Edge, NJ, 1991.
- [Bul70] V. K. Bulitko. Equations and inequalities in a free group and a free semigroup. *Tul. Gos. Ped. Inst. Učen. Zap. Mat. Kaf.*, (2 Geometr. i Algebra):242–252, 1970.
- [BW01] Martin R. Bridson and Daniel T. Wise. Malnormality is undecidable in hyperbolic groups. *Israel J. Math.*, 124:313–316, 2001.
- [Can84] James W. Cannon. The combinatorial structure of cocompact discrete hyperbolic groups. *Geom. Dedicata*, 16(2):123–148, 1984.
- [CDP90] M. Coornaert, T. Delzant, and A. Papadopoulos. *Géométrie et théorie des groupes*, volume 1441 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1990. Les groupes hyperboliques de Gromov. [Gromov hyperbolic groups], With an English summary.
- [Cha95] Olivier Chapuis. Universal theory of certain solvable groups and bounded Ore group rings. *J. Algebra*, 176(2):368–391, 1995.

- [Cul84] Marc Culler. Finite groups of outer automorphisms of a free group. In *Contributions to group theory*, volume 33 of *Contemp. Math.*, pages 197–207. Amer. Math. Soc., Providence, RI, 1984.
- [Dah05] Francois Dahmani. Existential questions in (relatively) hyperbolic groups, 2005. *Israel Journal of Math.* 173 (2009) 91–124.
- [Del95] T. Delzant. L’image d’un groupe dans un groupe hyperbolique. *Comment. Math. Helv.*, 70(2):267–284, 1995.
- [DG] François Dahmani and Vincent Guirardel. The isomorphism problem for all hyperbolic groups. arXiv:1002.2590 [math.GR].
- [DG08] François Dahmani and Daniel Groves. The isomorphism problem for toral relatively hyperbolic groups. *Publ. Math. Inst. Hautes Études Sci.*, (107):211–290, 2008.
- [DGH05] Volker Diekert, Claudio Gutierrez, and Christian Hagenah. The existential theory of equations with rational constraints in free groups is PSPACE-complete. *Inform. and Comput.*, 202(2):105–140, 2005.
- [Die02] Volker Diekert. Makanin’s algorithm. In M. Lothaire, editor, *Algebraic combinatorics on words*, volume 90 of *Encyclopedia of Mathematics and its Applications*, chapter 12, pages 387–442. Cambridge University Press, 2002.
- [DL08] Volker Diekert and Markus Lohrey. Word equations over graph products. *Internat. J. Algebra Comput.*, 18(3):493–533, 2008.
- [DM06] Volker Diekert and Anca Muscholl. Solvability of equations in graph groups is decidable. *Internat. J. Algebra Comput.*, 16(6):1047–1069, 2006.
- [Gab97] Damien Gaboriau. Générateurs indépendants pour les systèmes d’isométries de dimension un. *Ann. Inst. Fourier (Grenoble)*, 47(1):101–122, 1997.
- [GLP94] D. Gaboriau, G. Levitt, and F. Paulin. Pseudogroups of isometries of $\mathbb{R}$ and Rips’ theorem on free actions on $\mathbb{R}$ -trees. *Israel J. Math.*, 87(1-3):403–428, 1994.
- [GLP95] Damien Gaboriau, Gilbert Levitt, and Frédéric Paulin. Pseudogroups of isometries of $\mathbb{R}$: reconstruction of free actions on $\mathbb{R}$ -trees. *Ergodic Theory Dynam. Systems*, 15(4):633–652, 1995.
- [Gro87] M. Gromov. Hyperbolic groups. In *Essays in group theory*, volume 8 of *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New York, 1987.
- [Gui98] Vincent Guirardel. Approximations of stable actions on $\mathbb{R}$ -trees. *Comment. Math. Helv.*, 73(1):89–121, 1998.
- [Gus00] Paulo Gusmão. Feuilletages mesurés et pseudogroupes d’isométries du cercle. *J. Math. Sci. Univ. Tokyo*, 7(3):487–508, 2000.
- [Kap96] Ilya Kapovich. Detecting quasiconvexity: algorithmic aspects. *Geometric and computational perspectives on infinite groups (Minneapolis, MN and New Brunswick, NJ, 1994)*, 91–99, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 25, Amer. Math. Soc., Providence, RI, 1996.
- [KM05] Olga Kharlampovich and Alexei Myasnikov. Implicit function theorem over free groups. *J. Algebra*, 290(1):1–203, 2005.

- [KM06] Olga Kharlampovich and Alexei Myasnikov. Elementary theory of free non-abelian groups. *J. Algebra*, 302(2):451–552, 2006.
- [KP98] Antoni Kościelski and Leszek Pacholski. Makanin’s algorithm is not primitive recursive. *Theoret. Comput. Sci.*, 191(1-2):145–156, 1998.
- [LS06a] Markus Lohrey and Géraud Senizergues. Equations in hnn extensions, 2006. short version in proceedings ICALP’06, p. 504-515.
- [LS06b] Markus Lohrey and Géraud Senizergues. Rational subsets of hnn extensions, 2006. short version in proceedings ICALP’06, LNCS 4052, p. 504-515.
- [Mak82] G. S. Makanin. Equations in a free group. *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(6):1199–1273, 1344, 1982.
- [Mak84] G. S. Makanin. Decidability of the universal and positive theories of a free group. *Izv. Akad. Nauk SSSR Ser. Mat.*, 48(4):735–749, 1984.
- [Mat70] Ju. V. Matiyasevich. The Diophantineness of enumerable sets. *Dokl. Akad. Nauk SSSR*, 191:279–282, 1970.
- [MK02] V. D. Mazurov and E. I. Khukhro, editors. *The Kourovka notebook*. Rossiiskaya Akademiya Nauk Sibirskoe Otdelenie, Institut Matematiki im. S. L. Soboleva, Novosibirsk, augmented edition, 2002. Unsolved problems in group theory.
- [Pap96] P. Papasoglu. An algorithm detecting hyperbolicity. In *Geometric and computational perspectives on infinite groups (Minneapolis, MN and New Brunswick, NJ, 1994)*, volume 25 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 193–200. Amer. Math. Soc., Providence, RI, 1996.
- [Pla75] J. F. Plante. Measure preserving pseudogroups and a theorem of Sacksteder. *Ann. Inst. Fourier (Grenoble)*, 25(1):vii, 237–249, 1975.
- [Pla04] Wojciech Plandowski. Satisfiability of word equations with constants is in PSPACE. *J. ACM*, 51(3):483–496 (electronic), 2004.
- [Raz84] A. A. Razborov. Systems of equations in a free group. *Izv. Akad. Nauk SSSR Ser. Mat.*, 48(4):779–832, 1984.
- [Rei] Cornelius Reinfeldt. In preparation.
- [Rip82] E. Rips. Subgroups of small cancellation groups. *Bull. London Math. Soc.*, 14(1):45–47, 1982.
- [Rom79a] V. A. Roman’kov. Equations in free metabelian groups. *Sibirsk. Mat. Zh.*, 20(3):671–673, 694, 1979.
- [Rom79b] V. A. Roman’kov. Universal theory of nilpotent groups. *Mat. Zametki*, 25(4):487–495, 635, 1979.
- [RS95] E. Rips and Z. Sela. Canonical representatives and equations in hyperbolic groups. *Invent. Math.*, 120(3):489–512, 1995.
- [Sch86] Alexander Schrijver. *Theory of linear and integer programming*. Wiley-Interscience Series in Discrete Mathematics. John Wiley & Sons Ltd., Chichester, 1986. A Wiley-Interscience Publication.

- [Sch92] Klaus U. Schulz. Makanin’s algorithm for word equations—two improvements and a generalization. In *Word equations and related topics (Tübingen, 1990)*, volume 572 of *Lecture Notes in Comput. Sci.*, pages 85–150. Springer, Berlin, 1992.
- [Sel95] Z. Sela. The isomorphism problem for hyperbolic groups. I. *Ann. of Math. (2)*, 141(2):217–283, 1995.
- [Sel01] Zlil Sela. Diophantine geometry over groups. I. Makanin-Razborov diagrams. *Publ. Math. Inst. Hautes Études Sci.*, 93:31–105, 2001.
- [Sel06] Z. Sela. Diophantine geometry over groups. VI. The elementary theory of a free group. *Geom. Funct. Anal.*, 16(3):707–730, 2006.
- [Tru95] J. K. Truss. Equation-solving in free nilpotent groups of class 2 and 3. *Bull. London Math. Soc.*, 27(1):39–45, 1995.
- [Zim81] Bruno Zimmermann. Über Homöomorphismen n -dimensionaler Henkelkörper und endliche Erweiterungen von Schottky-Gruppen. *Comment. Math. Helv.*, 56(3):474–486, 1981.

François Dahmani, Vincent Guirardel
 Institut de Mathématiques de Toulouse (UMR 5219)
 Université Paul Sabatier, Toulouse III
 31062 Toulouse cedex 9.
 France.
e-mail: dahmani@math.univ-toulouse.fr, guirardel@math.univ-toulouse.fr